



JABchem



Not to be shared without the copyright holder's permission










Past Papers Advanced Higher Chemistry

2023 Marking Scheme













Grade Awarded	Mark Required		% candidates achieving grade
	(/120)	%	
A	77+	64.2%	%
B	62+	51.7%	%
C	48+	40.0%	%
D	33+	27.5%	%
No award	<33	<27.5%	%

Section:	Multiple Choice	Extended Answer	Project
Average Mark:	15.7 /25	50.4 /95	No Project in 2023

2023 Adv Higher Chemistry Marking Scheme

MC Qu	Answer	Reasoning																																																
1	D	<input checked="" type="checkbox"/> A Aufbau principle states that 4s fills before 3d shells <input checked="" type="checkbox"/> B Aufbau principle states that 4s fills before 3d shells <input checked="" type="checkbox"/> C 4s ¹ only occurs when 3d is either half-filled 3d ⁵ in Cr or completely full 3d ¹⁰ in Cu <input checked="" type="checkbox"/> D 4s is filled before 3d																																																
2	C	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <thead> <tr> <th colspan="2">Linear</th> <th colspan="2">Tetrahedral</th> <th colspan="2">Trigonal Pyramidal</th> <th colspan="2">Trigonal Bipyramidal</th> </tr> <tr> <td colspan="2">2</td> <td colspan="2">4</td> <td colspan="2">4</td> <td colspan="2">5</td> </tr> <tr> <td colspan="2">electron pairs</td> <td colspan="2">electron pairs</td> <td colspan="2">electron pairs</td> <td colspan="2">electron pairs</td> </tr> <tr> <td>2</td> <td>0</td> <td>4</td> <td>0</td> <td>3</td> <td>1</td> <td>5</td> <td>0</td> </tr> <tr> <td>bonding pairs</td> <td>lone pairs</td> <td>bonding pairs</td> <td>lone pairs</td> <td>bonding pairs</td> <td>lone pairs</td> <td>bonding pairs</td> <td>lone pairs</td> </tr> </thead> <tbody> <tr> <td colspan="2">Cl — Be — Cl</td> <td colspan="2">  </td> <td colspan="2">  </td> <td colspan="2">  </td> </tr> </tbody> </table>	Linear		Tetrahedral		Trigonal Pyramidal		Trigonal Bipyramidal		2		4		4		5		electron pairs		electron pairs		electron pairs		electron pairs		2	0	4	0	3	1	5	0	bonding pairs	lone pairs	bonding pairs	lone pairs	bonding pairs	lone pairs	bonding pairs	lone pairs	Cl — Be — Cl							
Linear		Tetrahedral		Trigonal Pyramidal		Trigonal Bipyramidal																																												
2		4		4		5																																												
electron pairs		electron pairs		electron pairs		electron pairs																																												
2	0	4	0	3	1	5	0																																											
bonding pairs	lone pairs	bonding pairs	lone pairs	bonding pairs	lone pairs	bonding pairs	lone pairs																																											
Cl — Be — Cl																																																		
3	B	<input checked="" type="checkbox"/> A s-block found in groups 1+2 <input checked="" type="checkbox"/> B p-block found in groups 3→0 <input checked="" type="checkbox"/> C d-block found in transition metals between groups 2+3 <input checked="" type="checkbox"/> D f-block found in bottom 2 rows of Periodic Table																																																
4	A	<input checked="" type="checkbox"/> A Heterogeneous catalysis work by adsorption of reactants onto active sites of catalyst and homogeneous catalyst are in same state as catalysts. <input checked="" type="checkbox"/> B an incomplete d-subshell is required for the interaction of the catalyst with reactant(s) <input checked="" type="checkbox"/> C heterogeneous catalyst is in a different state to the reactants <input checked="" type="checkbox"/> D adsorption onto active sites is the mechanism of heterogeneous catalysts																																																
5	D	Halfway point of steep vertical region (neutralisation point) of line is approx. pH=6 <input checked="" type="checkbox"/> A acidic pH at start of experiment ∴ alkali is being added to an acid <input checked="" type="checkbox"/> B strong alkali/weak acid gives alkaline pH on neutralisation but pH is acidic at neutralisation <input checked="" type="checkbox"/> C acidic pH at start of experiment ∴ alkali is being added to an acid <input checked="" type="checkbox"/> D pH at neutralisation is acidic weak alkali and strong acid gives acidic pH on neutralisation																																																
6	D	<table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse; text-align: center;"> <thead> <tr> <th>Term</th> <th>Bronsted-Lowry Definition</th> </tr> </thead> <tbody> <tr> <td>acid</td> <td>Species which donates H⁺ and becomes conjugate base</td> </tr> <tr> <td>base</td> <td>Species which accepts H⁺ and becomes conjugate acid</td> </tr> <tr> <td>conjugate acid</td> <td>Species formed when base accepted an H⁺ ion</td> </tr> <tr> <td>conjugate base</td> <td>Species formed when acid donated an H⁺ ion</td> </tr> </tbody> </table> <p>H₃O⁺ ions being absorbed by conjugate base turning back into acid Weak acid donates H₃O⁺ ions and becomes conjugate base in the process.</p>	Term	Bronsted-Lowry Definition	acid	Species which donates H ⁺ and becomes conjugate base	base	Species which accepts H ⁺ and becomes conjugate acid	conjugate acid	Species formed when base accepted an H ⁺ ion	conjugate base	Species formed when acid donated an H ⁺ ion																																						
Term	Bronsted-Lowry Definition																																																	
acid	Species which donates H ⁺ and becomes conjugate base																																																	
base	Species which accepts H ⁺ and becomes conjugate acid																																																	
conjugate acid	Species formed when base accepted an H ⁺ ion																																																	
conjugate base	Species formed when acid donated an H ⁺ ion																																																	
7	A	<input checked="" type="checkbox"/> A reactions which are feasible have negative values of ΔG and equilibrium favours products <input checked="" type="checkbox"/> B a feasible reaction is one that tends towards the products rather than the reactants <input checked="" type="checkbox"/> C a reaction with a positive value for standard free energy (ΔG) is not feasible <input checked="" type="checkbox"/> D a reaction with a positive value for standard free energy (ΔG) is not feasible																																																
8	B	<table style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td style="text-align: center;">①</td> <td style="text-align: center;">$\frac{1}{2}\text{N}_2 + \text{O}_2 \rightarrow \text{NO}_2$</td> <td style="text-align: right;">$\Delta G^\circ = +51.8 \text{ kJ mol}^{-1}$</td> </tr> <tr> <td style="text-align: center;">②</td> <td style="text-align: center;">$\text{N}_2 + 2\text{O}_2 \rightarrow \text{N}_2\text{O}_4$</td> <td style="text-align: right;">$\Delta G^\circ = +97.7 \text{ kJ mol}^{-1}$</td> </tr> <tr> <td style="text-align: center;">① x-2</td> <td style="text-align: center;">$2\text{NO}_2 \rightarrow \text{N}_2 + 2\text{O}_2$</td> <td style="text-align: right;">$\Delta G^\circ = -103.6 \text{ kJ mol}^{-1}$</td> </tr> <tr> <td style="text-align: center;">②</td> <td style="text-align: center;">$\text{N}_2 + 2\text{O}_2 \rightarrow \text{N}_2\text{O}_4$</td> <td style="text-align: right;">$\Delta G^\circ = +97.7 \text{ kJ mol}^{-1}$</td> </tr> <tr> <td style="text-align: center;">cancel out</td> <td style="text-align: center;"> $2\text{NO}_2 \rightarrow \cancel{\text{N}_2} + \cancel{2\text{O}_2}$ $\cancel{\text{N}_2} + \cancel{2\text{O}_2} \rightarrow \text{N}_2\text{O}_4$ </td> <td style="text-align: right;">$\Delta G^\circ = -5.9 \text{ kJ mol}^{-1}$</td> </tr> <tr> <td style="text-align: center;">Final equation</td> <td style="text-align: center;">$2\text{NO}_2 \rightarrow \text{N}_2\text{O}_4$</td> <td style="text-align: right;">$\Delta G^\circ = -5.9 \text{ kJ mol}^{-1}$</td> </tr> </tbody> </table>	①	$\frac{1}{2}\text{N}_2 + \text{O}_2 \rightarrow \text{NO}_2$	$\Delta G^\circ = +51.8 \text{ kJ mol}^{-1}$	②	$\text{N}_2 + 2\text{O}_2 \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = +97.7 \text{ kJ mol}^{-1}$	① x-2	$2\text{NO}_2 \rightarrow \text{N}_2 + 2\text{O}_2$	$\Delta G^\circ = -103.6 \text{ kJ mol}^{-1}$	②	$\text{N}_2 + 2\text{O}_2 \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = +97.7 \text{ kJ mol}^{-1}$	cancel out	$2\text{NO}_2 \rightarrow \cancel{\text{N}_2} + \cancel{2\text{O}_2}$ $\cancel{\text{N}_2} + \cancel{2\text{O}_2} \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = -5.9 \text{ kJ mol}^{-1}$	Final equation	$2\text{NO}_2 \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = -5.9 \text{ kJ mol}^{-1}$																														
①	$\frac{1}{2}\text{N}_2 + \text{O}_2 \rightarrow \text{NO}_2$	$\Delta G^\circ = +51.8 \text{ kJ mol}^{-1}$																																																
②	$\text{N}_2 + 2\text{O}_2 \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = +97.7 \text{ kJ mol}^{-1}$																																																
① x-2	$2\text{NO}_2 \rightarrow \text{N}_2 + 2\text{O}_2$	$\Delta G^\circ = -103.6 \text{ kJ mol}^{-1}$																																																
②	$\text{N}_2 + 2\text{O}_2 \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = +97.7 \text{ kJ mol}^{-1}$																																																
cancel out	$2\text{NO}_2 \rightarrow \cancel{\text{N}_2} + \cancel{2\text{O}_2}$ $\cancel{\text{N}_2} + \cancel{2\text{O}_2} \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = -5.9 \text{ kJ mol}^{-1}$																																																
Final equation	$2\text{NO}_2 \rightarrow \text{N}_2\text{O}_4$	$\Delta G^\circ = -5.9 \text{ kJ mol}^{-1}$																																																

9	C	<p>Positive ΔS value means an increase in disorder.</p> <p><input checked="" type="checkbox"/> A burning solid carbon into the gas CO_2 is an increase in disorder and exothermic</p> <p><input checked="" type="checkbox"/> B water vapour turning into solid snowflakes is a decrease in disorder and exothermic</p> <p><input checked="" type="checkbox"/> C liquid ethoxyethane (ether) evaporating into gas is endothermic and increases disorder</p> <p><input checked="" type="checkbox"/> D two gases becoming one solid is a decrease in disorder and exothermic</p>
10	A	<p>Rate = $k[X][Y]$ means that both X and Y are first order and 1 molecule of each appears as reactants in the slow rate determining step. $\therefore X + Y \rightarrow$ intermediate</p> <p>The overall reaction is $X + 2Y \rightarrow Z$ so there must be a second step where the intermediate formed in (slow) rate determining step reacts with a further molecule of Y.</p>
11	A	<p>The single carbon - carbon bond in alkanes is a sigma bond. Sigma bonds are formed by the end-on overlap of atomic orbitals along the axis of the covalent bond.</p> <p>In a non-polar covalent bond in an alkane, the bonding molecular orbital is symmetrical about the midpoint between two atoms.</p> <p>(Polar covalent bonds result from bonding molecular orbitals that are asymmetric about the midpoint between two atoms.)</p>
12	C	<p><input checked="" type="checkbox"/> A blue-green light is absorbed and not transmitted. Only red colour is transmitted.</p> <p><input checked="" type="checkbox"/> B electrons move from HOMO to LUMO to absorb a wavelength of light</p> <p><input checked="" type="checkbox"/> C electrons move from HOMO to LUMO and blue-green light absorbed and red transmitted</p> <p><input checked="" type="checkbox"/> D electrons move from HOMO to LUMO to absorb a wavelength of light</p>
13	D	<p><input checked="" type="checkbox"/> A ethanol contains O-H bonds and has hydrogen bonding between molecules</p> <p><input checked="" type="checkbox"/> B ethylamine contains N-H bonds and has hydrogen bonding between molecules</p> <p><input checked="" type="checkbox"/> C ethanoic acid contains O-H bonds and has hydrogen bonding between molecules</p> <p><input checked="" type="checkbox"/> D ethoxyethane has no N-H, O-H or H-F bonds and no hydrogen bonding in pure substance</p>
14	A	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> $\begin{array}{c} \text{H} \quad \text{Br} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$ <p>secondary bromoalkane</p> </div> <div style="text-align: center;"> <p>nucleophilic substitution</p> \longrightarrow <p>aqueous sodium hydroxide</p> </div> <div style="text-align: center;"> $\begin{array}{c} \text{H} \quad \text{OH} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$ <p>secondary alcohol</p> </div> <div style="text-align: center;"> <p>oxidation</p> \longrightarrow <p>acidified dichromate solution</p> </div> <div style="text-align: center;"> $\begin{array}{c} \text{H} \quad \text{O} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array}$ <p>Ketone</p> </div> </div>
15	C	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \quad \text{O} \\ \quad \quad \quad // \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C} \\ \quad \quad \quad \backslash \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$ <p>2-methylbutanal</p> </div> <div style="text-align: center;"> <p>lithium aluminium hydride</p> \longrightarrow <p>reduction</p> </div> <div style="text-align: center;"> $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{OH} \\ \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$ <p>2-methylbutan-1-ol</p> </div> </div> <ul style="list-style-type: none"> $CH_3CH_2CH(CH_3)CHO$ is an aldehyde called 2-methylbutanal $LiAlH_4$ is the reducing agent lithium aluminium hydride Aldehydes are reduced to primary alcohols by $LiAlH_4$
16	B	$C_4H_8O_2 + NH_3 \rightarrow X \rightarrow Y + H_2O$ $C_4H_8O_2 + NH_3 \rightarrow C_4H_{11}NO_2 \rightarrow C_4H_9NO + H_2O$ <p><input checked="" type="checkbox"/> A $CH_3CH_2CH_2CN$ has a molecular formula of C_4H_7N</p> <p><input checked="" type="checkbox"/> B $CH_3CH_2CH_2CONH_2$ has a molecular formula of C_4H_9NO</p> <p><input checked="" type="checkbox"/> C $CH_3CH_2CH_2CH_2NH_2$ has a molecular formula of $C_4H_{11}N$</p> <p><input checked="" type="checkbox"/> D $CH_3CH_2CH_2COONH_2$ has a molecular formula of $C_4H_9NO_2$</p>

17	A	<p>Methoxypropane has the structure: $\begin{array}{ccccccc} & & & & \text{H} & & \text{H} & \text{H} & \text{H} \\ & & & & & & & & \\ \text{H} & - & \text{C} & - & \text{O} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{H} \\ & & & & & & & & \\ & & & & \text{H} & & \text{H} & \text{H} & \text{H} \end{array}$</p> <p>It has a molecular formula of $\text{C}_4\text{H}_{10}\text{O}$</p> <p><input checked="" type="checkbox"/> A butanone $\text{CH}_3\text{CH}_2\text{COCH}_3$ has a molecular formula of $\text{C}_4\text{H}_8\text{O}$ <input checked="" type="checkbox"/> B ethoxyethane $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ has a molecular formula of $\text{C}_4\text{H}_{10}\text{O}$ <input checked="" type="checkbox"/> C butan-2-ol $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$ has a molecular formula of $\text{C}_4\text{H}_{10}\text{O}$ <input checked="" type="checkbox"/> D 2-methylpropan-1-ol $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{OH}$ has a molecular formula of $\text{C}_4\text{H}_{10}\text{O}$</p>																
18	B	<table border="1"> <thead> <tr> <th data-bbox="331 510 620 566">A</th> <th data-bbox="620 510 914 566">B</th> <th data-bbox="914 510 1208 566">C</th> <th data-bbox="1208 510 1500 566">D</th> </tr> </thead> <tbody> <tr> <td data-bbox="331 566 620 645">CH_2CHBr</td> <td data-bbox="620 566 914 645">CHClCHCH_3</td> <td data-bbox="914 566 1208 645">$\text{CH}_3\text{CH}_2\text{CHCl}_2$</td> <td data-bbox="1208 566 1500 645">$\text{CH}_3\text{C}(\text{CH}_3)\text{CHCH}_3$</td> </tr> <tr> <td data-bbox="331 645 620 869">  <p style="text-align: center;">bromoethane</p> </td> <td data-bbox="620 645 914 869">  <p style="text-align: center;">trans-1-chloropropene</p> </td> <td data-bbox="914 645 1208 869">  <p style="text-align: center;">1,1-dichlorobut-1-ene</p> </td> <td data-bbox="1208 645 1500 869">  <p style="text-align: center;">2-methylbut-2-ene</p> </td> </tr> <tr> <td data-bbox="331 869 620 1037">Left end of C=C double bond has two H groups attached. This prevents it from being a geometric isomer</td> <td data-bbox="620 869 914 1037">There are two (non-identical) groups, one on each end of the C=C double bond. This makes it a geometric isomer</td> <td data-bbox="914 869 1208 1037">Left end of C=C double bond has two H groups attached. This prevents it from being a geometric isomer</td> <td data-bbox="1208 869 1500 1037">Left end of C=C double bond has two CH₃ groups attached. This prevents it from being a geometric isomer</td> </tr> </tbody> </table>	A	B	C	D	CH_2CHBr	CHClCHCH_3	$\text{CH}_3\text{CH}_2\text{CHCl}_2$	$\text{CH}_3\text{C}(\text{CH}_3)\text{CHCH}_3$	 <p style="text-align: center;">bromoethane</p>	 <p style="text-align: center;">trans-1-chloropropene</p>	 <p style="text-align: center;">1,1-dichlorobut-1-ene</p>	 <p style="text-align: center;">2-methylbut-2-ene</p>	Left end of C=C double bond has two H groups attached. This prevents it from being a geometric isomer	There are two (non-identical) groups, one on each end of the C=C double bond. This makes it a geometric isomer	Left end of C=C double bond has two H groups attached. This prevents it from being a geometric isomer	Left end of C=C double bond has two CH ₃ groups attached. This prevents it from being a geometric isomer
A	B	C	D															
CH_2CHBr	CHClCHCH_3	$\text{CH}_3\text{CH}_2\text{CHCl}_2$	$\text{CH}_3\text{C}(\text{CH}_3)\text{CHCH}_3$															
 <p style="text-align: center;">bromoethane</p>	 <p style="text-align: center;">trans-1-chloropropene</p>	 <p style="text-align: center;">1,1-dichlorobut-1-ene</p>	 <p style="text-align: center;">2-methylbut-2-ene</p>															
Left end of C=C double bond has two H groups attached. This prevents it from being a geometric isomer	There are two (non-identical) groups, one on each end of the C=C double bond. This makes it a geometric isomer	Left end of C=C double bond has two H groups attached. This prevents it from being a geometric isomer	Left end of C=C double bond has two CH ₃ groups attached. This prevents it from being a geometric isomer															
19	A	<p><input checked="" type="checkbox"/> A in mass spectroscopy, the molecule breaks up during the process to give different lines <input checked="" type="checkbox"/> B in infrared spectroscopy, molecule has certain bonds stretched at certain wavenumbers <input checked="" type="checkbox"/> C in melting point analysis, a pure substance will melt first but will freeze back again. <input checked="" type="checkbox"/> D Proton NMR uses radio waves to flip nuclei but does not destroy the sample.</p>																
20	D	<table border="1"> <thead> <tr> <th data-bbox="331 1198 555 1232">Type of Drug</th> <th data-bbox="555 1198 815 1232">Binding Site</th> <th data-bbox="815 1198 1493 1232">Response</th> </tr> </thead> <tbody> <tr> <td data-bbox="331 1232 555 1276">Agonist</td> <td data-bbox="555 1232 815 1276">receptor binding site</td> <td data-bbox="815 1232 1493 1276">produces a response similar to the body's natural response</td> </tr> <tr> <td data-bbox="331 1276 555 1321">Antagonist</td> <td data-bbox="555 1276 815 1321">receptor binding site</td> <td data-bbox="815 1276 1493 1321">blocks the body's natural response</td> </tr> <tr> <td data-bbox="331 1321 555 1379">Enzyme Inhibitor</td> <td data-bbox="555 1321 815 1379">enzyme active site</td> <td data-bbox="815 1321 1493 1379">blocks the action of the enzyme</td> </tr> </tbody> </table>	Type of Drug	Binding Site	Response	Agonist	receptor binding site	produces a response similar to the body's natural response	Antagonist	receptor binding site	blocks the body's natural response	Enzyme Inhibitor	enzyme active site	blocks the action of the enzyme				
Type of Drug	Binding Site	Response																
Agonist	receptor binding site	produces a response similar to the body's natural response																
Antagonist	receptor binding site	blocks the body's natural response																
Enzyme Inhibitor	enzyme active site	blocks the action of the enzyme																
21	D	<p><input checked="" type="checkbox"/> A As doxepin is only molecule with O atom at top, this part not in common fragment <input checked="" type="checkbox"/> B Molecule B is incorrect at bottom the nitrogen part is rotatory and will flip round <input checked="" type="checkbox"/> C Doxepin lacks the -Cl group on top left of molecule ∴ Cl not in common fragment <input checked="" type="checkbox"/> D Molecule D is correct as the nitrogen group is rotary & flips round and found in all drugs</p>																
22	B	<table border="1"> <tbody> <tr> <td data-bbox="708 1541 900 1603">Element</td> <td data-bbox="900 1541 1011 1603">V</td> <td data-bbox="1011 1541 1123 1603">O</td> </tr> <tr> <td data-bbox="708 1603 900 1666">Mass or %</td> <td data-bbox="900 1603 1011 1666">61.4%</td> <td data-bbox="1011 1603 1123 1666">38.6</td> </tr> <tr> <td data-bbox="708 1666 900 1809">No. of moles <small>(divide % by gfm)</small></td> <td data-bbox="900 1666 1011 1809"> $\frac{61.4}{50.9}$ = 1.21 </td> <td data-bbox="1011 1666 1123 1809"> $\frac{38.6}{16}$ = 2.41 </td> </tr> <tr> <td data-bbox="708 1809 900 1980">Mole ratio <small>(divide through by smallest value)</small></td> <td data-bbox="900 1809 1011 1980"> $\frac{1.21}{1.21}$ = 1 </td> <td data-bbox="1011 1809 1123 1980"> $\frac{2.41}{1.21}$ = 1.99 = 2 </td> </tr> <tr> <td data-bbox="708 1980 900 2031">Empirical Formula</td> <td colspan="2" data-bbox="900 1980 1123 2031">VO₂</td> </tr> </tbody> </table>	Element	V	O	Mass or %	61.4%	38.6	No. of moles <small>(divide % by gfm)</small>	$\frac{61.4}{50.9}$ = 1.21	$\frac{38.6}{16}$ = 2.41	Mole ratio <small>(divide through by smallest value)</small>	$\frac{1.21}{1.21}$ = 1	$\frac{2.41}{1.21}$ = 1.99 = 2	Empirical Formula	VO ₂		
Element	V	O																
Mass or %	61.4%	38.6																
No. of moles <small>(divide % by gfm)</small>	$\frac{61.4}{50.9}$ = 1.21	$\frac{38.6}{16}$ = 2.41																
Mole ratio <small>(divide through by smallest value)</small>	$\frac{1.21}{1.21}$ = 1	$\frac{2.41}{1.21}$ = 1.99 = 2																
Empirical Formula	VO ₂																	



23	C	<p>A primary standard must:</p> <table border="1" data-bbox="389 143 1442 206"> <thead> <tr> <th data-bbox="389 143 651 206">be available in a high state of purity</th> <th data-bbox="651 143 916 206">be stable when solid and in solution</th> <th data-bbox="916 143 1177 206">be soluble</th> <th data-bbox="1177 143 1442 206">have a reasonably high GFM</th> </tr> </thead> </table> <p>Examples of primary standards include:</p> <table data-bbox="373 232 1458 367"> <tr> <td data-bbox="373 232 555 295">sodium carbonate Na_2CO_3</td> <td data-bbox="555 232 868 295">hydrated oxalic acid $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$</td> <td data-bbox="868 232 1177 295">potassium hydrogen phthalate $\text{KH}(\text{C}_8\text{H}_4\text{O}_4)$</td> </tr> <tr> <td data-bbox="373 295 555 367">silver nitrate AgNO_3</td> <td data-bbox="555 295 868 367">potassium iodate KIO_3</td> <td data-bbox="868 295 1177 367">potassium dichromate $\text{K}_2\text{Cr}_2\text{O}_7$</td> </tr> </table> <p><input checked="" type="checkbox"/> A calcium carbonate is insoluble and therefore not a primary standard</p> <p><input checked="" type="checkbox"/> B Hydrochloric acid</p> <p><input checked="" type="checkbox"/> C sodium carbonate meets the four criteria listed in table above</p> <p><input checked="" type="checkbox"/> D Sodium hydroxide is not a primary standard as it has a relatively low GFM, is unstable as a solid (absorbs moisture) and unstable as a solution. Sodium hydroxide solution must be standardised before being used in volumetric analysis.</p>	be available in a high state of purity	be stable when solid and in solution	be soluble	have a reasonably high GFM	sodium carbonate Na_2CO_3	hydrated oxalic acid $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$	potassium hydrogen phthalate $\text{KH}(\text{C}_8\text{H}_4\text{O}_4)$	silver nitrate AgNO_3	potassium iodate KIO_3	potassium dichromate $\text{K}_2\text{Cr}_2\text{O}_7$
be available in a high state of purity	be stable when solid and in solution	be soluble	have a reasonably high GFM									
sodium carbonate Na_2CO_3	hydrated oxalic acid $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$	potassium hydrogen phthalate $\text{KH}(\text{C}_8\text{H}_4\text{O}_4)$										
silver nitrate AgNO_3	potassium iodate KIO_3	potassium dichromate $\text{K}_2\text{Cr}_2\text{O}_7$										
24	B	<p style="text-align: center;">$420 \text{ ppm} = 420 \text{ mg CO}_2 \text{ per } 1 \text{ kg atmosphere}$</p> <p style="text-align: center;">$1 \text{ kg atmosphere} = 0.42 \text{ g CO}_2$</p> <p style="text-align: center;">$5.1 \times 10^{18} \text{ kg atmosphere} = 0.42 \text{ g CO}_2 \times 5.1 \times 10^{18}$</p> <p style="text-align: center;">$= 2.142 \times 10^{18} \text{ g CO}_2$</p> <p style="text-align: center;">$= 2.1 \times 10^{15} \text{ kg CO}_2$</p>										
25	B	$\begin{array}{ccccccc} \text{Pb}^{2+} & + & \text{CrO}_4^{2-} & \rightarrow & \text{PbCrO}_4 & & \\ 1\text{mol} & & & & 1\text{mol} & & \end{array}$ $\begin{array}{ccccccc} \text{PbCrO}_4 & + & 2\text{Cl}^- & \rightarrow & \text{PbCl}_2 & + & \text{CrO}_4^{2-} \\ 1\text{mol} & & & & & & 1\text{mol} \end{array}$ $\begin{array}{ccccccc} 2\text{CrO}_4^{2-} & + & 6\text{I}^- & + & 16\text{H}^+ & \rightarrow & 2\text{Cr}^{3+} & + & 3\text{I}_2 & + & 8\text{H}_2\text{O} \\ 2\text{mol} & & & & & & 3\text{mol} & & & & \\ 1\text{mol} & & & & & & 1.5\text{mol} & & & & \end{array}$										




2023 Adv Higher Chemistry Marking Scheme

Long Qu	Answer	Reasoning																																
1a(i)	Answer to include:	<table border="1"> <tr> <td>1st mark:</td> <td>(Excited) electrons fall to a lower energy level</td> </tr> <tr> <td>2nd mark:</td> <td>Difference in energy levels corresponds to $\left[\begin{array}{c} \text{red light} \\ \text{light} \\ \text{energy} \end{array} \right]$ emitted.</td> </tr> </table>	1 st mark:	(Excited) electrons fall to a lower energy level	2 nd mark:	Difference in energy levels corresponds to $\left[\begin{array}{c} \text{red light} \\ \text{light} \\ \text{energy} \end{array} \right]$ emitted.																												
1 st mark:	(Excited) electrons fall to a lower energy level																																	
2 nd mark:	Difference in energy levels corresponds to $\left[\begin{array}{c} \text{red light} \\ \text{light} \\ \text{energy} \end{array} \right]$ emitted.																																	
1a(ii)	<table border="1"> <tr> <td></td> <td></td> <td></td> <td>m_s</td> </tr> <tr> <td>1</td> <td></td> <td></td> <td></td> </tr> </table>				m_s	1				<p>Li has electronic configuration of $1s^2 2s^1$. Li^+ has electronic configuration of $1s^2$</p> <table border="1"> <tr> <td>n</td> <td>1s subshell is in 1st shell and has a value for Principal quantum number (n) of n=1</td> </tr> <tr> <td>l</td> <td>s subshells have an angular momentum quantum number (l) of l=0</td> </tr> <tr> <td>m_l</td> <td>s subshells have a magnetic quantum number (m_l) of -l to +l \therefore 0</td> </tr> <tr> <td>m_s</td> <td>All electrons either have a spin magnetic quantum number of either $+\frac{1}{2}$ or $-\frac{1}{2}$</td> </tr> </table>	n	1s subshell is in 1 st shell and has a value for Principal quantum number (n) of n=1	l	s subshells have an angular momentum quantum number (l) of l=0	m_l	s subshells have a magnetic quantum number (m_l) of -l to +l \therefore 0	m_s	All electrons either have a spin magnetic quantum number of either $+\frac{1}{2}$ or $-\frac{1}{2}$																
			m_s																															
1																																		
n	1s subshell is in 1 st shell and has a value for Principal quantum number (n) of n=1																																	
l	s subshells have an angular momentum quantum number (l) of l=0																																	
m_l	s subshells have a magnetic quantum number (m_l) of -l to +l \therefore 0																																	
m_s	All electrons either have a spin magnetic quantum number of either $+\frac{1}{2}$ or $-\frac{1}{2}$																																	
1a(iii)A	+7 or VII	Oxidation state of O = -2 \therefore 4xO in ClO_4^- combine to equal -8. Perchlorate ClO_4^- has charge of -1 over whole ion \therefore the oxidation state of Cl must be +7																																
1a(iii)B	9	<table border="1"> <thead> <tr> <th>Bond</th> <th>Number of bonds</th> <th>Number of sigma</th> <th>Number of Pi bonds</th> </tr> </thead> <tbody> <tr> <td>N - H</td> <td>3</td> <td>$3 \times 1 = 3$</td> <td>0</td> </tr> <tr> <td>N - C</td> <td>1</td> <td>1</td> <td>0</td> </tr> <tr> <td>C = N</td> <td>1</td> <td>1</td> <td>1</td> </tr> <tr> <td>C - N</td> <td>1</td> <td>1</td> <td>0</td> </tr> <tr> <td>N - N</td> <td>2</td> <td>$2 \times 1 = 2$</td> <td>0</td> </tr> <tr> <td>N = N</td> <td>1</td> <td>1</td> <td>1</td> </tr> <tr> <td>Total</td> <td></td> <td>9</td> <td>2</td> </tr> </tbody> </table>	Bond	Number of bonds	Number of sigma	Number of Pi bonds	N - H	3	$3 \times 1 = 3$	0	N - C	1	1	0	C = N	1	1	1	C - N	1	1	0	N - N	2	$2 \times 1 = 2$	0	N = N	1	1	1	Total		9	2
Bond	Number of bonds	Number of sigma	Number of Pi bonds																															
N - H	3	$3 \times 1 = 3$	0																															
N - C	1	1	0																															
C = N	1	1	1																															
C - N	1	1	0																															
N - N	2	$2 \times 1 = 2$	0																															
N = N	1	1	1																															
Total		9	2																															
1b(i)A	-1648	$\begin{aligned} \Delta H^\circ &= \Sigma \Delta H_f^\circ(\text{products}) - \Sigma \Delta H_f^\circ(\text{reactants}) \\ &= (2 \times -824) - (4 \times 0) + (3 \times 0) \\ &= -1648 - 0 \\ &= -1648 \text{ kJ mol}^{-1} \end{aligned}$																																
1b(i)B	-549.4	$\begin{aligned} \Delta S^\circ &= \Sigma S^\circ(\text{products}) - \Sigma S^\circ(\text{reactants}) \\ &= (2 \times 87.4) - (4 \times 27.3) + (3 \times 205) \\ &= 174.8 - (109.2 + 615) \\ &= 174.8 - 724.2 \\ &= -549.4 \text{ J K}^{-1} \text{ mol}^{-1} \end{aligned}$																																
1b(i)C	3000	$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ = 0 \quad \therefore T = \frac{\Delta H^\circ}{\Delta S^\circ} = \frac{-1648 \times 1000 \text{ J mol}^{-1}}{-549.4 \text{ J K}^{-1} \text{ mol}^{-1}} = 3000\text{K}$																																
1b(ii)	2694g	<p>gfm $NaClO_3 = (1 \times 23) + (1 \times 35.5) + (3 \times 16) = 23 + 35.5 + 48 = 106.5\text{g}$</p> <p>1 person in 1 minute uses 0.380 litres oxygen 5 persons in 1 minute uses 0.380 litres oxygen \times 5 $= 1.9$ litres oxygen</p> <p>5 persons in 1 hour uses 1.9 litres oxygen \times 60 $= 114$ litres oxygen</p> <p>5 persons in 8 hours uses 114 litres oxygen \times 8 $= 912$ litres oxygen</p> <p>36 litres oxygen produced from 1 mol $NaClO_3$ 912 litres oxygen produced from 1 mol $NaClO_3 \times \frac{912}{36}$ $= 25.3$ mol $NaClO_3$</p> <p>mass = no. of mol \times gfm = $25.3 \times 106.5 = 2694\text{g}$</p>																																
2a(i)	1 st Order	<table border="1"> <thead> <tr> <th>Experiments</th> <th>$[HgCl_2]$</th> <th>$[C_2O_4^{2-}]$</th> <th>Effect of Rate</th> <th>Order</th> </tr> </thead> <tbody> <tr> <td>3+2</td> <td>$\times 2$</td> <td>same</td> <td>$\times 2$</td> <td>$[HgCl_2]^1$</td> </tr> </tbody> </table>	Experiments	$[HgCl_2]$	$[C_2O_4^{2-}]$	Effect of Rate	Order	3+2	$\times 2$	same	$\times 2$	$[HgCl_2]^1$																						
Experiments	$[HgCl_2]$	$[C_2O_4^{2-}]$	Effect of Rate	Order																														
3+2	$\times 2$	same	$\times 2$	$[HgCl_2]^1$																														
2a(ii)	2 nd Order	<table border="1"> <thead> <tr> <th>Experiments</th> <th>$[HgCl_2]$</th> <th>$[C_2O_4^{2-}]$</th> <th>Effect of Rate</th> <th>Order</th> </tr> </thead> <tbody> <tr> <td>1+2</td> <td>same</td> <td>$\times 2$</td> <td>$\times 4$</td> <td>$[C_2O_4^{2-}]^2$</td> </tr> </tbody> </table>	Experiments	$[HgCl_2]$	$[C_2O_4^{2-}]$	Effect of Rate	Order	1+2	same	$\times 2$	$\times 4$	$[C_2O_4^{2-}]^2$																						
Experiments	$[HgCl_2]$	$[C_2O_4^{2-}]$	Effect of Rate	Order																														
1+2	same	$\times 2$	$\times 4$	$[C_2O_4^{2-}]^2$																														






2b	rate = $k[\text{HgCl}_2][\text{C}_2\text{O}_4^{2-}]^2$	rate = $k \times [\text{HgCl}_2]^1 \times [\text{C}_2\text{O}_4^{2-}]^2 = k[\text{HgCl}_2][\text{C}_2\text{O}_4^{2-}]^2$						
2c(i)	$2.56 \times 10^{-4} \text{ l}^2 \text{ mol}^{-2} \text{ s}^{-1}$ Or $0.000256 \text{ l}^2 \text{ mol}^{-2} \text{ s}^{-1}$	Using Experiment 1 (although experiments 2+3 can also be used) $k = \frac{\text{rate}}{[\text{HgCl}_2][\text{C}_2\text{O}_4^{2-}]^2} = \frac{0.860 \times 10^{-6} \text{ mol l}^{-1} \text{ s}^{-1}}{0.0840 \text{ mol l}^{-1} \times (0.200)^2 \text{ mol}^2 \text{ l}^{-2}} = 2.56 \times 10^{-4} \text{ l}^2 \text{ mol}^{-2} \text{ s}^{-1}$						
2c(ii)	0.508	For experiment 4: Rate = $k \times [\text{HgCl}_2] \times [\text{C}_2\text{O}_4^{2-}]^2$ $2.11 \times 10^{-6} = 2.56 \times 10^{-4} \times 0.0320 \times [\text{C}_2\text{O}_4^{2-}]^2$ $[\text{C}_2\text{O}_4^{2-}]^2 = \frac{2.11 \times 10^{-6}}{2.56 \times 10^{-4} \times 0.0320} = 0.258 \text{ mol}^2 \text{ l}^{-2}$ $[\text{C}_2\text{O}_4^{2-}] = \sqrt{0.258} \text{ mol l}^{-1} = 0.508 \text{ mol l}^{-1}$						
3a(i)	Add more/excess calcium chloride solution (to see if further precipitation occurs)	When precipitation appears to be complete, the mixture is allowed to settle and more reactant is added to see if there is any further precipitation/cloudiness showing the reaction is incomplete and further reactant is added. If there is no more precipitation/cloudiness then the precipitate can be removed by vacuum filtration.						
3a(ii)	The precipitate should be dried/heated in an oven and weighed until constant mass	When precipitate is removed from the filtrate by vacuum filtration, some of the water remains in the precipitate residue. Heating the precipitate in an oven will drive off water. When the precipitate is removed from the oven it is cooled in a desiccator and the mass measured in a balance. Heating in oven, cooling in a desiccator and weight on balance is repeated until constant mass is achieved.						
3b	0.59	gfm $\text{Ca}(\text{COO})_2 = (1 \times 40.1) + (2 \times 12) + (4 \times 16) = 40.1 + 24 + 64 = 128.1 \text{ g}$ $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{0.075}{128.1} = 5.85 \times 10^{-4} \text{ mol}$ $(\text{COOH})_2 + \text{CaCl}_2 \longrightarrow \text{Ca}(\text{COO})_2 + 2\text{HCl}$ $\begin{matrix} 1 \text{ mol} & & 1 \text{ mol} \\ 5.85 \times 10^{-4} \text{ mol} & & 5.85 \times 10^{-4} \text{ mol} \end{matrix}$ gfm $(\text{COOH})_2 = (2 \times 12) + (4 \times 16) + (2 \times 1) = 24 + 64 + 2 = 90 \text{ g}$ $\text{mass} = \text{no. of mol} \times \text{gfm} = 5.85 \times 10^{-4} \times 90 = 0.0527 \text{ g}$ $\% \text{ mass} = \frac{\text{mass of oxalic acid}}{\text{mass of leaves}} \times 100 = \frac{0.0527}{8.975} \times 100 = 0.59\%$						
3c	One answer from:	<table border="1"> <tbody> <tr> <td>Not all oxalic acid extracted out</td> <td>Not soaked in water for long enough</td> <td>Not all precipitated out/not gone to completion</td> </tr> <tr> <td>Not enough CaCl_2 added</td> <td>Transfer loss</td> <td>Natural variation in samples</td> </tr> </tbody> </table>	Not all oxalic acid extracted out	Not soaked in water for long enough	Not all precipitated out/not gone to completion	Not enough CaCl_2 added	Transfer loss	Natural variation in samples
Not all oxalic acid extracted out	Not soaked in water for long enough	Not all precipitated out/not gone to completion						
Not enough CaCl_2 added	Transfer loss	Natural variation in samples						
4	Open Question to include:	<table border="1"> <thead> <tr> <th>3 mark answer</th> <th>2 mark answer</th> <th>1 mark answer</th> </tr> </thead> <tbody> <tr> <td>Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.</td> <td>Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.</td> <td>Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.</td> </tr> </tbody> </table>	3 mark answer	2 mark answer	1 mark answer	Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
3 mark answer	2 mark answer	1 mark answer						
Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.						
5a	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^8$	Charge in Nickel ion = +2 as H_2O is a neutral ligand so +2 charge on complex comes from Ni^{2+} ion. $\text{Ni atom: } 1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2$ 4s electrons are removed before 3d electrons						


		Ni^{2+} ion: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8$																												
5b	Hexaamminenickel(II)	<p>Hexaamminenickel(II) = $[\text{Ni}(\text{NH}_3)_6]^{2+}$</p> <p>no. of ligands NH_3 ligand metal name charge on metal ion</p> <table border="1"> <tr> <td colspan="2">Neutral ligands include:</td> <td colspan="2">Negative Ligands include:</td> <td>Central Ion:</td> <td>Charge:</td> </tr> <tr> <td>Ligand</td> <td>Name</td> <td>Ligand</td> <td>Name</td> <td>Positive Complex: metals keep their name</td> <td rowspan="3">Charge of central ion is converted into roman numerals and put in brackets</td> </tr> <tr> <td>H_2O</td> <td>aqua</td> <td>Chloride Cl^-</td> <td>chlorido</td> <td>Negative Complex: Metals end in ATE</td> </tr> <tr> <td>NH_3</td> <td>ammine</td> <td>Cyanide CN^-</td> <td>cyanido</td> <td>e.g. Cuprate, Ferrate, Cobaltate</td> </tr> <tr> <td></td> <td>CO</td> <td>carbonyl</td> <td>Nitrite NO_2^-</td> <td>nitrito</td> <td></td> </tr> </table>	Neutral ligands include:		Negative Ligands include:		Central Ion:	Charge:	Ligand	Name	Ligand	Name	Positive Complex: metals keep their name	Charge of central ion is converted into roman numerals and put in brackets	H_2O	aqua	Chloride Cl^-	chlorido	Negative Complex: Metals end in ATE	NH_3	ammine	Cyanide CN^-	cyanido	e.g. Cuprate, Ferrate, Cobaltate		CO	carbonyl	Nitrite NO_2^-	nitrito	
Neutral ligands include:		Negative Ligands include:		Central Ion:	Charge:																									
Ligand	Name	Ligand	Name	Positive Complex: metals keep their name	Charge of central ion is converted into roman numerals and put in brackets																									
H_2O	aqua	Chloride Cl^-	chlorido	Negative Complex: Metals end in ATE																										
NH_3	ammine	Cyanide CN^-	cyanido	e.g. Cuprate, Ferrate, Cobaltate																										
	CO	carbonyl	Nitrite NO_2^-	nitrito																										
5c(i)	Bidentate	A bidentate ligand has two parts of the ligand that can donate pairs of electrons to the central metal ion at the centre of the complex.																												
5c(ii)	6	6 different pairs of electrons across three bidentate ligands are donated to central metal ion. This gives a co-ordination number of 6.																												
5d(i)	Answer to include:	<table border="1"> <tr> <td>1st mark:</td> <td>Electrons move to higher energy d orbitals</td> </tr> <tr> <td>2nd mark:</td> <td> <p>purple light 380-400nm light red and blue light complementary colour (to green)</p> <p>} absorbed</p> </td> </tr> </table>	1 st mark:	Electrons move to higher energy d orbitals	2 nd mark:	<p>purple light 380-400nm light red and blue light complementary colour (to green)</p> <p>} absorbed</p>																								
1 st mark:	Electrons move to higher energy d orbitals																													
2 nd mark:	<p>purple light 380-400nm light red and blue light complementary colour (to green)</p> <p>} absorbed</p>																													
5d(ii)	One answer from:	<p>Ammonia absorbs</p> <ul style="list-style-type: none"> shorter wavelength higher energy higher frequency uv 																												
6a(i)	$K = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]}$	For reaction: $aA + bB \rightleftharpoons cC + dD$ $K = \frac{[\text{C}]^c [\text{D}]^d}{[\text{A}]^a [\text{B}]^b}$																												
6a(ii)	981.8	<p>I_2 at start = 0.25mol I_2 at equilibrium = 0.015mol</p> <p>$\therefore \text{I}_2$ reacted = 0.25mol - 0.015mol = 0.235mol</p> <table style="margin-left: auto; margin-right: auto;"> <tr> <td>H_2</td> <td>+</td> <td>I_2</td> <td>\rightleftharpoons</td> <td>2HI</td> </tr> <tr> <td>1mol</td> <td></td> <td>1mol</td> <td></td> <td>2mol</td> </tr> <tr> <td>0.235mol</td> <td></td> <td>0.235mol</td> <td></td> <td>0.470mol</td> </tr> <tr> <td>(reacted)</td> <td></td> <td>(reacted)</td> <td></td> <td>(formed)</td> </tr> </table> <p>H_2 remaining at equilibrium = 0.25mol - 0.235mol = 0.015mol</p> $K = \frac{[\text{HI}]^2}{[\text{H}_2][\text{I}_2]} = \frac{(0.470)^2}{0.015 \times 0.015} = 981.8$	H_2	+	I_2	\rightleftharpoons	2HI	1mol		1mol		2mol	0.235mol		0.235mol		0.470mol	(reacted)		(reacted)		(formed)								
H_2	+	I_2	\rightleftharpoons	2HI																										
1mol		1mol		2mol																										
0.235mol		0.235mol		0.470mol																										
(reacted)		(reacted)		(formed)																										
6a(iii)	207.2	$E = \frac{L \times h \times c}{\lambda} = \frac{6.02 \times 10^{23} \text{ mol}^{-1} \times 6.63 \times 10^{-34} \text{ J s} \times 3 \times 10^8 \text{ m s}^{-1}}{578 \times 10^{-9} \text{ m}}$ $= 207159 \text{ J mol}^{-1}$ $= 207.2 \text{ kJ mol}^{-1}$																												
6b(i)	Buchner Funnel or Hirsch Funnel or Sintered Glass Funnel	The Buchner funnel allows vacuum filtration to place which speeds up the flow of filtrate through the Buchner funnel. The reduced pressure provides a faster means to separate precipitate from filtrate																												

6b(ii)	87.4%	<p>gfm $I_2 = (2 \times 126.9) = 253.8g$</p> $\text{no. of mol } I_2 = \frac{\text{mass}}{\text{gfm}} = \frac{285}{253.8} = 1.12 \text{ mol}$ $I_2 + H_2S \longrightarrow 2HI + S$ <p style="text-align: center;"> 1mol 2mol </p> <p style="text-align: center;"> 1.12mol 2.25mol (theoretical) </p> <p>gfm $HI = (1 \times 1) + (1 \times 126.9) = 1 + 126.9 = 127.9g$</p> $\text{no. of mol } HI = \frac{\text{mass}}{\text{gfm}} = \frac{251}{127.9} = 1.96 \text{ mol (actual)}$ $\% \text{ Yield} = \frac{\text{actual}}{\text{theoretical}} \times 100 = \frac{1.96}{2.25} \times 100 = 87.4\%$																					
6c		<p>Notes on example:</p> <ul style="list-style-type: none"> Hydroxyl -OH groups become carbonyl C=O groups C-C single bond between carbons with hydroxyl -OH groups splits 																					
7a	Concentration of dissolved CO_2 decreases as equilibrium moves to left.	<p>Increase in temperature favours the endothermic reaction</p> <p>Reverse reaction is favoured as reverse reaction is endothermic</p> <p>Reverse reaction decreases $H_2CO_{3(aq)}$ concentration and releases $CO_{2(g)}$ into air.</p>																					
7b(i)	<p>Carbonic acid is a weaker acid as it has higher pK_a / lower K_a or</p> <p>Ethanoic acid is a stronger acid as it has lower pK_a / higher K_a</p>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 20%;">Weak Acid</th> <th style="width: 30%;">K_a</th> <th style="width: 50%;">pK_a</th> </tr> </thead> <tbody> <tr> <td>Carbonic acid</td> <td style="text-align: center;">4.5×10^{-7}</td> <td style="text-align: center;">6.35</td> </tr> <tr> <td>Ethanoic acid</td> <td style="text-align: center;">1.7×10^{-5}</td> <td style="text-align: center;">4.76</td> </tr> <tr> <td>Conclusion</td> <td>Carbonic acid is a lower strength acid/weaker acid as it has a smaller value of K_a</td> <td>Carbonic acid is a lower strength acid/weaker acid as it has a higher value of pK_a</td> </tr> </tbody> </table>	Weak Acid	K _a	pK _a	Carbonic acid	4.5×10^{-7}	6.35	Ethanoic acid	1.7×10^{-5}	4.76	Conclusion	Carbonic acid is a lower strength acid/weaker acid as it has a smaller value of K _a	Carbonic acid is a lower strength acid/weaker acid as it has a higher value of pK _a									
Weak Acid	K _a	pK _a																					
Carbonic acid	4.5×10^{-7}	6.35																					
Ethanoic acid	1.7×10^{-5}	4.76																					
Conclusion	Carbonic acid is a lower strength acid/weaker acid as it has a smaller value of K _a	Carbonic acid is a lower strength acid/weaker acid as it has a higher value of pK _a																					
7b(ii)	<p style="text-align: center;">Base (or proton acceptor)</p>	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%; text-align: center;">$H_2CO_{3(aq)}$</td> <td style="width: 5%; text-align: center;">+</td> <td style="width: 25%; text-align: center;">$H_2O(l)$</td> <td style="width: 5%; text-align: center;">\rightleftharpoons</td> <td style="width: 25%; text-align: center;">$H_3O^+(aq)$</td> <td style="width: 5%; text-align: center;">+</td> <td style="width: 20%; text-align: center;">$HCO_3^-(aq)$</td> </tr> <tr> <td style="text-align: center;">Acid</td> <td></td> <td style="text-align: center;">Base</td> <td></td> <td style="text-align: center;">Conjugate Acid</td> <td></td> <td style="text-align: center;">Conjugate base</td> </tr> <tr> <td style="text-align: center;">loses a proton/H^+ (proton donor)</td> <td></td> <td style="text-align: center;">gains a proton/H^+ (proton acceptor)</td> <td></td> <td style="text-align: center;">formed when base accepts a proton/H^+</td> <td></td> <td style="text-align: center;">formed when acid loses a proton/H^+</td> </tr> </table>	$H_2CO_{3(aq)}$	+	$H_2O(l)$	\rightleftharpoons	$H_3O^+(aq)$	+	$HCO_3^-(aq)$	Acid		Base		Conjugate Acid		Conjugate base	loses a proton/ H^+ (proton donor)		gains a proton/ H^+ (proton acceptor)		formed when base accepts a proton/ H^+		formed when acid loses a proton/ H^+
$H_2CO_{3(aq)}$	+	$H_2O(l)$	\rightleftharpoons	$H_3O^+(aq)$	+	$HCO_3^-(aq)$																	
Acid		Base		Conjugate Acid		Conjugate base																	
loses a proton/ H^+ (proton donor)		gains a proton/ H^+ (proton acceptor)		formed when base accepts a proton/ H^+		formed when acid loses a proton/ H^+																	
7c(i)	6.31×10^{-9}	<p style="text-align: center;">pH = 8.2</p> <p style="text-align: center;">$-\log_{10}[H_3O^+] = 8.2$</p> <p style="text-align: center;">$\log_{10}[H_3O^+] = -8.2$</p> <p style="text-align: center;">$[H_3O^+] = 10^{-8.2}$</p> <p style="text-align: center;">$[H_3O^+] = 6.31 \times 10^{-9} \text{ mol l}^{-1}$</p>																					

7c(ii)	99.7%	$\text{pH} = 7.9$ $-\log_{10}[\text{H}_3\text{O}^+] = 7.9$ $\log_{10}[\text{H}_3\text{O}^+] = -7.9$ $[\text{H}_3\text{O}^+] = 10^{-7.9}$ $[\text{H}_3\text{O}^+] = 1.26 \times 10^{-8} \text{ mol l}^{-1}$	$\% \text{ Increase} = \frac{\text{difference in value}}{\text{original value}} \times 100$ $\% \text{ Increase} = \frac{1.26 \times 10^{-8} - 6.31 \times 10^{-9}}{6.31 \times 10^{-9}} \times 100$ $\% \text{ Increase} = \frac{6.29 \times 10^{-9}}{6.31 \times 10^{-9}} \times 100$ $\% \text{ Increase} = 99.7\%$																		
7d(i)	Answer to include:	Reacting known mass of seashells with an excess volume of a hydrochloric acid of known concentration	The resulting mixture is then titrated with sodium hydroxide to work out the number of moles of hydrochloric acid in excess	From the initial number of moles of hydrochloric acid, the number of moles used in the reaction can be determined.	The initial number of moles of the calcium carbonate in seashells being analysed can then be calculated.																
7d(i)	Answer to include: All 4 points = 2marks 2 or 3 points = 1 mark	React calcium carbonate /shells with HCl/acid	Known quantity/moles/volume and concentration of HCl/acid	idea of excess HCl/acid	titrate with NaOH.																
7d(ii)	Repeat using pure calcium carbonate	In a control experiment, a known mass of pure calcium carbonate can assess the accuracy of the reaction method.																			
8a(i)	15-crown-5	In example given 18-crown-6: 18 refers to the number of atoms in the ring and 6 refers to number of oxygens The question: 15 atoms in the ring and 5 oxygen atoms in ring \therefore 15-crown-5																			
8a(ii)	 Also accepted: 	<table border="1"> <thead> <tr> <th>Element</th> <th>Member of Group 1</th> <th>Number of O in ether</th> <th>Name</th> </tr> </thead> <tbody> <tr> <td>Li</td> <td>1st</td> <td>4</td> <td>12-crown-4</td> </tr> <tr> <td>Na</td> <td>2nd</td> <td>5</td> <td>15-crown-5</td> </tr> <tr> <td>K</td> <td>3rd</td> <td>6</td> <td>18-crown-6</td> </tr> </tbody> </table>				Element	Member of Group 1	Number of O in ether	Name	Li	1 st	4	12-crown-4	Na	2 nd	5	15-crown-5	K	3 rd	6	18-crown-6
Element	Member of Group 1	Number of O in ether	Name																		
Li	1 st	4	12-crown-4																		
Na	2 nd	5	15-crown-5																		
K	3 rd	6	18-crown-6																		
8a(iii)	Can donate lone (non-bonding) pairs of electrons	Ligands have lone pairs of electrons which form dative covalent bonds by donating pairs of electrons to the central metal ion forming dative covalent bonds. In crown ethers, the lone pair electrons are pointing into the centre of the ring where the metal ion is located.																			
8b(i)	$\text{C}_6\text{H}_{14}\text{O}_4$																				
8b(ii)	Condensation or nucleophilic substitution	H from one molecule and OH from next molecule are removed as the molecules join together forming H ₂ O																			
8c	2.43×10^{-3} or 0.0024	gfm 18-Crown-6 = 264g $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{0.225}{264} = 8.52 \times 10^{-4} \text{ mol}$ As K ⁺ ions and 18-crown-6 are in 1:1 ratio: $8.52 \times 10^{-4} \text{ mol K}^+$ ions $\text{concentration} = \frac{\text{no. of mol}}{\text{volume}} = \frac{8.52 \times 10^{-4} \text{ mol}}{0.350 \text{ litres}} = 2.43 \times 10^{-3} \text{ mol l}^{-1}$																			

9	Open Question to include:	3 mark answer	2 mark answer	1 mark answer
		Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
10a(i)	One answer from:	Adjacent (unhybridised) p orbitals that overlap/form a molecular orbital across a system/number of carbon atoms	Electrons delocalised across a number of carbon atoms	Alternating single and double bonds
10a(ii)	sp^2	C=C double bonds contain sp^2 hybridisation with a C-C sigma bond down the axis of the bond and side on overlap of unhybridised orbitals to form the pi bond.		
10a(iii)	One from:			
10b(i)	Repulsion from pi electrons or double bond	The electrons Cl-Cl bond are repelled by the pi electrons in the C=C double bond and this polarises the Cl-Cl bond forming a δ^+ end and a δ^- end		
10b(ii)				
10b(iii)A	Restricted rotation/ lack of free rotation (around the single bonds) in the ring	<p>The ring of six carbons prevents the rotation so there are two locked positions for side groups above and below the ring with no rotation.</p> <ul style="list-style-type: none"> if both side groups are above ring then this is the cis geometric isomer if one side group is above ring and the other side group below ring then this is the trans geometric isomer. <p>A C=C double bond also has a lack of rotation in the bond due to the Pi bond so that also has the geometric isomers</p>		

10b(iii)B	<p>structure drawn of 1,1-dichlorocyclohexane</p>  <p>or</p>  <p>1,4-dichlorocyclohexane</p>	<p>An optical isomer has a chiral carbon with four different groups attached.</p> <p>1,1-dichlorocyclohexane has no carbons with four different groups. C_1 has two chlorine atoms and the other carbons each have two hydrogens.</p> <p>1,4-dichlorocyclohexane lacks a chiral carbons as C_1 and C_4 each have a hydrogen atom and a chlorine atom but the ring of carbons is symmetrical and are identical regardless of which way you go around the ring.</p>									
10c(i)	<p>Diagram showing:</p> <p>1 mark for curly arrows 1 mark for correct intermediate</p>										
10c(ii)	1.08	<p>$e = 1.60 \times 10^{-19} C$ partial charge = 0.178</p> <p>$Q = \text{partial charge} \times e = 0.178 \times 1.60 \times 10^{-19} C = 2.848 \times 10^{-20} C$</p> <p>$Q = 2.848 \times 10^{-20} C$ $r = 0.127 \text{ nm} = 0.127 \times 10^{-9} \text{ m} = 1.27 \times 10^{-10} \text{ m}$</p> <p>$\mu = Q \times r = 2.848 \times 10^{-20} C \times 1.27 \times 10^{-10} \text{ m} = 3.62 \times 10^{-30} \text{ Cm}$</p> <p>$3.34 \times 10^{-30} \text{ Cm} = 1 \text{ D}$</p> <p>$3.62 \times 10^{-30} \text{ Cm} = 1 \text{ D} \times \frac{3.62 \times 10^{-30} \text{ Cm}}{3.34 \times 10^{-30} \text{ Cm}}$</p> <p>$= 1.08 \text{ D}$</p>									
11a(i)	NO_2^+	<p>Nitration electrophilic substitution reactions involve concentrated nitric acid and concentrated sulphuric acid and forms the electrophile nitronium NO_2^+ ion</p> <p>$\text{HNO}_3 + \text{H}_2\text{SO}_4 \longrightarrow \text{NO}_2^+ + \text{H}_3\text{O}^+ + \text{HSO}_4^-$</p> <p style="text-align: center;"> <small>concentrated nitric acid concentrated sulphuric acid nitronium ion hydronium ion hydrogensulphate ion</small> </p>									
11a(ii)A											
11a(ii)B											
11b	Reduction	<p style="text-align: center;">Decrease in the oxygen : hydrogen ratio</p> <table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>Formula</th> <th>$\text{C}_6\text{H}_4\text{O}_3$</th> <th>$\text{C}_6\text{H}_7\text{O}$</th> </tr> </thead> <tbody> <tr> <td>O : H ratio</td> <td>3 : 4</td> <td>1 : 7</td> </tr> <tr> <td></td> <td>1 : 1.33</td> <td></td> </tr> </tbody> </table>	Formula	$\text{C}_6\text{H}_4\text{O}_3$	$\text{C}_6\text{H}_7\text{O}$	O : H ratio	3 : 4	1 : 7		1 : 1.33	
Formula	$\text{C}_6\text{H}_4\text{O}_3$	$\text{C}_6\text{H}_7\text{O}$									
O : H ratio	3 : 4	1 : 7									
	1 : 1.33										

11c	Ethanoic acid or Ethanoic anhydride or Ethanoyl chloride	Amines react with carboxylic acids to form alkylammonium salts and form an amide when heated.																																									
11d	Answer to include:	2 marks for all 4 points in table.			1 mark for 2 or 3 points in table																																						
		minimum/small volume (of solvent)	hot water/solvent	cool	filter																																						
11e(i)A	C=O or Carbonyl																																										
11e(i)B	Overlap of peaks from different bonds with one example as shown:	O-H peak in same region as $\begin{cases} \text{N-H} \\ \text{C-H} \end{cases}$ or N-H peak in same region as $\begin{cases} \text{O-H} \\ \text{C-H} \end{cases}$																																									
11e(ii)A	<p>CH₃-C=O circled:</p> 	<table border="1" data-bbox="576 600 1495 748"> <tr> <td>Chemical Shift (ppm)</td> <td>2.1</td> <td>6.7</td> <td>7.3</td> <td>9.1</td> <td>9.8</td> </tr> <tr> <td>Height of Curve (mm)</td> <td>50</td> <td>33</td> <td>33</td> <td>17</td> <td>17</td> </tr> <tr> <td>No of Hydrogens in Group</td> <td>3</td> <td>2</td> <td>2</td> <td>1</td> <td>1</td> </tr> <tr> <td>Multiplet</td> <td>singlet</td> <td>doublet</td> <td>doublet</td> <td>singlet</td> <td>singlet</td> </tr> <tr> <td>No of Hydrogen next door</td> <td>0</td> <td>2</td> <td></td> <td>0</td> <td>0</td> </tr> <tr> <td>Group</td> <td>CH₃C=O</td> <td>ArH?</td> <td></td> <td>-HN-C=O</td> <td>Ar-OH</td> </tr> </table>						Chemical Shift (ppm)	2.1	6.7	7.3	9.1	9.8	Height of Curve (mm)	50	33	33	17	17	No of Hydrogens in Group	3	2	2	1	1	Multiplet	singlet	doublet	doublet	singlet	singlet	No of Hydrogen next door	0	2		0	0	Group	CH ₃ C=O	ArH?		-HN-C=O	Ar-OH
Chemical Shift (ppm)	2.1	6.7	7.3	9.1	9.8																																						
Height of Curve (mm)	50	33	33	17	17																																						
No of Hydrogens in Group	3	2	2	1	1																																						
Multiplet	singlet	doublet	doublet	singlet	singlet																																						
No of Hydrogen next door	0	2		0	0																																						
Group	CH ₃ C=O	ArH?		-HN-C=O	Ar-OH																																						
11e(ii)B	<table border="1" data-bbox="343 763 496 853"> <tr> <td>Doublet</td> </tr> <tr> <td>Doublet</td> </tr> </table>	Doublet	Doublet																																								
Doublet																																											
Doublet																																											
11e(iii)A	Sample is pure with no extra peaks or only 5 (hydrogen) peaks	The same number of peaks with the same chemical shift indicated the sample is pure as contaminants would add other peaks to the NMR spectrum as they would contain other chemical groups.																																									
11e(iii)B	Mixed melting point or TLC or Chromatography	The melting point of the recrystallised paracetamol is measured using melting point apparatus. Pure paracetamol is added to see if the melting point is the same. If the melting point alters then the sample is not pure.																																									