



JABchem



Not to be shared without the copyright holder's permission

Past Papers Advanced Higher Chemistry




2019 Marking Scheme

Grade Awarded	Mark Required		% candidates achieving grade
	(/130)	%	
A	90+	69.2%	33.8%
B	77+	59.2%	26.3%
C	64+	49.2%	22.2%
D	57+	43.8%	7.3%
No award	<57	<43.8%	10.4%

Section:	Multiple Choice	Extended Answer	Assignment
Average Mark:	21.0 /30	40.4 /70	19.2 /30

2019 Adv Higher Chemistry Marking Scheme

MC Qu	Answer	% Pupils Correct	Reasoning								
1	B	76	<input checked="" type="checkbox"/> A adsorption spectra formed by adsorption of energy as electrons are promoted <input checked="" type="checkbox"/> B emission spectra are formed from the release of energy as electrons drop down <input checked="" type="checkbox"/> C 310nm is a wavelength in the ultraviolet not visible region of EM Spectrum <input checked="" type="checkbox"/> D 310nm is a wavelength in the ultraviolet not visible region of EM Spectrum								
2	D	85	In gravimetric analysis, heating $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$ will release the 10 water molecules trapped in the structure so that just Na_2CO_3 remains. The substance is heated, cooled in a desiccator and its mass measured. The process is repeated until the mass is constant.								
3	C	37	<input checked="" type="checkbox"/> A adding sodium nitrate to magnesium ions does not produce a magnesium precipitate <input checked="" type="checkbox"/> B adding silver(I) nitrate to magnesium ions does not produce a magnesium precipitate <input checked="" type="checkbox"/> C sodium carbonate forms a magnesium carbonate precipitate with magnesium ion solution <input checked="" type="checkbox"/> D silver(I) carbonate is insoluble so carbonate ions cannot precipitate with magnesium ions								
4	A	90	<input checked="" type="checkbox"/> A Hund's Rule: electrons occupy degenerate orbitals singly with parallel spins before pairing <input checked="" type="checkbox"/> B Pauli Exclusion Principle: no two electrons in an atom can have the same set of four quantum numbers <input checked="" type="checkbox"/> C Aufbau Principle: electrons occupy orbitals in order of increasing energy <input checked="" type="checkbox"/> D the energy of an electron in an atom is quantised into specific energy levels/electron shells								
5	D	75	<input checked="" type="checkbox"/> A BeCl_2 is a linear molecule with 180° angles <input checked="" type="checkbox"/> B BCl_3 is a trigonal planar molecule with 120° angles between bonds <input checked="" type="checkbox"/> C CCl_4 is a tetrahedral molecule with 109.5° angles between bonds <input checked="" type="checkbox"/> D PCl_5 is a trigonal pyramidal molecule with 90° (and 120°) angles between bonds								
6	D	51	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 25%;">Species</th> <th style="width: 25%;">Fe</th> <th style="width: 25%;">Fe²⁺</th> <th style="width: 25%;">Fe³⁺</th> </tr> </thead> <tbody> <tr> <td>Electronic configuration</td> <td>$1s^2 2s^2 2p^6 3s^2 2p^6 3d^6 4s^2$</td> <td>$1s^2 2s^2 2p^6 3s^2 2p^6 3d^6$</td> <td>$1s^2 2s^2 2p^6 3s^2 2p^6 3d^5$</td> </tr> </tbody> </table> <input checked="" type="checkbox"/> A Fe ²⁺ and Fe ³⁺ both have three occupied energy levels (n=1, n=2 and n=3) <input checked="" type="checkbox"/> B Fe ²⁺ has four unpaired electrons in 3d and Fe ³⁺ has five unpaired electrons in 3d <input checked="" type="checkbox"/> C Fe ²⁺ is higher in ECS than Fe ³⁺ . Fe ³⁺ is better oxidising agent than reducing agent <input checked="" type="checkbox"/> D Fe ³⁺ more stable because Fe ³⁺ had 5 unpaired electrons in half-filled d-subshell	Species	Fe	Fe ²⁺	Fe ³⁺	Electronic configuration	$1s^2 2s^2 2p^6 3s^2 2p^6 3d^6 4s^2$	$1s^2 2s^2 2p^6 3s^2 2p^6 3d^6$	$1s^2 2s^2 2p^6 3s^2 2p^6 3d^5$
Species	Fe	Fe ²⁺	Fe ³⁺								
Electronic configuration	$1s^2 2s^2 2p^6 3s^2 2p^6 3d^6 4s^2$	$1s^2 2s^2 2p^6 3s^2 2p^6 3d^6$	$1s^2 2s^2 2p^6 3s^2 2p^6 3d^5$								
7	B	71	<input checked="" type="checkbox"/> A $\text{Cr}_2\text{O}_7^{2-}$ ions have Cr with an oxidation state of +6 <input checked="" type="checkbox"/> B MnO_4^- ions have Mn with an oxidation state of +7 <input checked="" type="checkbox"/> C VO^{2+} ions have V with an oxidation state of +4 <input checked="" type="checkbox"/> D Sn^{4+} ions have Sn with an oxidation state of +4								
8	D	60	<input checked="" type="checkbox"/> A Co-ordination number of the copper 4 as there are four bonds from ligand to Cu <input checked="" type="checkbox"/> B Co-ordination number of the copper 4 as there are four bonds from ligand to Cu <input checked="" type="checkbox"/> C The ligand shares four pairs of electrons with the one metal ion = tetradentate <input checked="" type="checkbox"/> D Four bonds between ligand and central metal ion: co-ordination number equals 4 and is tetradentate								
9	A	72	<input checked="" type="checkbox"/> A Decrease in temperature favours the forward exothermic reaction. This gives more products and increasing the numerator on top of the equilibrium constant calculation which increases the value of equilibrium constant. <input checked="" type="checkbox"/> B Increase in temperature favours the reverse endothermic reaction. Lowers K value <input checked="" type="checkbox"/> C Changes in pressure do not alter the equilibrium constant <input checked="" type="checkbox"/> D Changes in pressure do not alter the equilibrium constant								
10	C	72	<input checked="" type="checkbox"/> A The reaction is feasible when the value of ΔG is below zero <input checked="" type="checkbox"/> B The reaction is only feasible when the value of ΔG is below zero (above 300K) <input checked="" type="checkbox"/> C The value of ΔG is negative above 300K so reaction is feasible above 300K <input checked="" type="checkbox"/> D The value of ΔG is positive below 300K so reaction is <u>not</u> feasible below 300K								
11	C	89	$\begin{aligned} \Delta H^\circ &= \Sigma \Delta H_f^\circ(\text{products}) - \Sigma \Delta H_f^\circ(\text{reactants}) \\ &= (2 \times 0) + (3 \times -242) - (1 \times -822) + (3 \times 0) \\ &= (0 - 726) - (-822 + 0) \\ &= -726 - (-822) \\ &= +96 \text{ kJ mol}^{-1} \end{aligned}$								

12	B	39	<input checked="" type="checkbox"/> A Steam condensing into water gives off heat $\therefore \Delta H$ is negative <input checked="" type="checkbox"/> B ΔH is negative as heat is given off. ΔS is negative as molecules are more ordered <input checked="" type="checkbox"/> C Steam condensing into water gives off heat $\therefore \Delta H$ is negative <input checked="" type="checkbox"/> D Molecules are closer together and more ordered in water $\therefore \Delta S$ is negative																		
13	A	86	<table border="1"> <thead> <tr> <th>Experiment</th> <th>Change</th> <th>Effect on Rate</th> <th>Order of reactant</th> </tr> </thead> <tbody> <tr> <td>1+2</td> <td>$[X] \times 2$</td> <td>$\times 2$</td> <td>$[X]^1$</td> </tr> <tr> <td>1+2</td> <td>$[Y] \times 2$</td> <td>No effect</td> <td>$[Y]^0$</td> </tr> </tbody> </table> $\text{Rate} = k [X]^1 [Y]^0 = k[X]$	Experiment	Change	Effect on Rate	Order of reactant	1+2	$[X] \times 2$	$\times 2$	$[X]^1$	1+2	$[Y] \times 2$	No effect	$[Y]^0$						
Experiment	Change	Effect on Rate	Order of reactant																		
1+2	$[X] \times 2$	$\times 2$	$[X]^1$																		
1+2	$[Y] \times 2$	No effect	$[Y]^0$																		
14	B	75	<table> <tbody> <tr> <td>rate = $k [A]^1 [B]^1$</td> <td>rate = $k [C]^2$</td> </tr> <tr> <td>$\therefore k = \frac{\text{rate}}{[A]^1 [B]^1}$</td> <td>$\therefore k = \frac{\text{rate}}{[C]^2}$</td> </tr> <tr> <td>= $\frac{\text{mol l}^{-1} \text{s}^{-1}}{\text{mol l}^{-1} \times \text{mol l}^{-1}}$</td> <td>= $\frac{\text{mol l}^{-1} \text{s}^{-1}}{\text{mol}^2 \text{l}^{-2}}$</td> </tr> <tr> <td>= $\text{l mol}^{-1} \text{s}^{-1}$</td> <td>= $\text{l mol}^{-1} \text{s}^{-1}$</td> </tr> </tbody> </table>	rate = $k [A]^1 [B]^1$	rate = $k [C]^2$	$\therefore k = \frac{\text{rate}}{[A]^1 [B]^1}$	$\therefore k = \frac{\text{rate}}{[C]^2}$	= $\frac{\text{mol l}^{-1} \text{s}^{-1}}{\text{mol l}^{-1} \times \text{mol l}^{-1}}$	= $\frac{\text{mol l}^{-1} \text{s}^{-1}}{\text{mol}^2 \text{l}^{-2}}$	= $\text{l mol}^{-1} \text{s}^{-1}$	= $\text{l mol}^{-1} \text{s}^{-1}$										
rate = $k [A]^1 [B]^1$	rate = $k [C]^2$																				
$\therefore k = \frac{\text{rate}}{[A]^1 [B]^1}$	$\therefore k = \frac{\text{rate}}{[C]^2}$																				
= $\frac{\text{mol l}^{-1} \text{s}^{-1}}{\text{mol l}^{-1} \times \text{mol l}^{-1}}$	= $\frac{\text{mol l}^{-1} \text{s}^{-1}}{\text{mol}^2 \text{l}^{-2}}$																				
= $\text{l mol}^{-1} \text{s}^{-1}$	= $\text{l mol}^{-1} \text{s}^{-1}$																				
15	C	78	<table border="1"> <thead> <tr> <th>Bonds</th> <th>2x C=C</th> <th>2x C-C</th> <th>1x C≡N</th> <th>2x C-O</th> <th>3x C-H</th> </tr> </thead> <tbody> <tr> <td>Sigma σ bonds</td> <td>2</td> <td>2</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>Pi π bonds</td> <td>2</td> <td>0</td> <td>2</td> <td>0</td> <td>0</td> </tr> </tbody> </table>	Bonds	2x C=C	2x C-C	1x C≡N	2x C-O	3x C-H	Sigma σ bonds	2	2	1	2	3	Pi π bonds	2	0	2	0	0
Bonds	2x C=C	2x C-C	1x C≡N	2x C-O	3x C-H																
Sigma σ bonds	2	2	1	2	3																
Pi π bonds	2	0	2	0	0																
16	A	89																			
17	C	49	<input checked="" type="checkbox"/> A The right C in C=C double bond has 2 -H groups \therefore no geometric isomerism <input checked="" type="checkbox"/> B The right C in C=C double bond has 2 -CH ₃ groups \therefore no geometric isomerism <input checked="" type="checkbox"/> C Both ends of C=C double bond have different groups \therefore has geometric isomerism <input checked="" type="checkbox"/> D The right C in C=C double bond has 2 -H groups \therefore no geometric isomerism																		
18	C	80	 <p>Structure: 3,4-dimethylhex-3-ene <i>Both ethyl groups are on same side of C=C double bond so this is the CIS geometric isomer.</i> Name: cis-3,4-dimethylhex-3-ene</p>																		
19	D	59	<input checked="" type="checkbox"/> A Trimethylamine is a tertiary amine so no hydrogen bonding \therefore lower boiling point <input checked="" type="checkbox"/> B Trimethylamine is a tertiary amine so no hydrogen bonding \therefore lower boiling point <input checked="" type="checkbox"/> C Trimethylamine has no H-N bonds so lacks any hydrogen bonding \therefore less soluble <input checked="" type="checkbox"/> D Trimethylamine has no hydrogen bonding as it lacks a H-N bond. This means trimethylamine molecules are further apart and this lowers the boiling point. The lack of N-H bonds lowers the solubility of trimethylamine in water.																		
20	B	78	Lithium Aluminium Hydride is a reducing agent for: <table style="margin-left: 40px;"> <tbody> <tr> <td>carboxylic acid</td> <td>\longrightarrow</td> <td>aldehyde</td> <td>\longrightarrow</td> <td>primary alcohol</td> </tr> <tr> <td>ketone</td> <td>\longrightarrow</td> <td>secondary alcohol</td> <td></td> <td></td> </tr> </tbody> </table>	carboxylic acid	\longrightarrow	aldehyde	\longrightarrow	primary alcohol	ketone	\longrightarrow	secondary alcohol										
carboxylic acid	\longrightarrow	aldehyde	\longrightarrow	primary alcohol																	
ketone	\longrightarrow	secondary alcohol																			
21	C	56	<table border="1"> <tbody> <tr> <td>Step 1</td> <td>Electrophilic substitution: -NO₂ group is substituted onto the benzene ring.</td> </tr> <tr> <td>Step 2</td> <td>Reduction: Decrease in the oxygen:hydrogen ratio as -NO₂ group is reduced to -NH₂ group</td> </tr> <tr> <td>Step 3</td> <td>Condensation: two molecules join together and small molecule removes where they join.</td> </tr> </tbody> </table>	Step 1	Electrophilic substitution: -NO ₂ group is substituted onto the benzene ring.	Step 2	Reduction: Decrease in the oxygen:hydrogen ratio as -NO ₂ group is reduced to -NH ₂ group	Step 3	Condensation: two molecules join together and small molecule removes where they join.												
Step 1	Electrophilic substitution: -NO ₂ group is substituted onto the benzene ring.																				
Step 2	Reduction: Decrease in the oxygen:hydrogen ratio as -NO ₂ group is reduced to -NH ₂ group																				
Step 3	Condensation: two molecules join together and small molecule removes where they join.																				
22	B	63																			
23	B	64	The heaviest m/z peak on a mass spectrum is the mass of the original compound. $\text{gfm } C_3H_6O = (3 \times 12) + (6 \times 1) + (1 \times 16) = 36 + 6 + 16 = 58\text{g}$																		




24	A	90	<table border="1"> <thead> <tr> <th>Element</th> <th>C</th> <th>H</th> <th>O</th> </tr> </thead> <tbody> <tr> <td>%</td> <td>80.0</td> <td>9.3</td> <td>10.7</td> </tr> <tr> <td>No. of moles (divide % by gfm)</td> <td>$\frac{80.0}{12}$ = 6.67</td> <td>$\frac{9.3}{1}$ = 9.300</td> <td>$\frac{10.7}{16}$ = 0.669</td> </tr> <tr> <td>Mole ratio (divide through by smallest value)</td> <td>$\frac{6.667}{0.669}$ = 9.96</td> <td>$\frac{9.300}{0.669}$ 13.90</td> <td>$\frac{0.669}{0.669}$ 1.00</td> </tr> <tr> <td>Round to Whole Number</td> <td>10</td> <td>14</td> <td>1</td> </tr> </tbody> </table>	Element	C	H	O	%	80.0	9.3	10.7	No. of moles (divide % by gfm)	$\frac{80.0}{12}$ = 6.67	$\frac{9.3}{1}$ = 9.300	$\frac{10.7}{16}$ = 0.669	Mole ratio (divide through by smallest value)	$\frac{6.667}{0.669}$ = 9.96	$\frac{9.300}{0.669}$ 13.90	$\frac{0.669}{0.669}$ 1.00	Round to Whole Number	10	14	1
			Element	C	H	O																	
			%	80.0	9.3	10.7																	
			No. of moles (divide % by gfm)	$\frac{80.0}{12}$ = 6.67	$\frac{9.3}{1}$ = 9.300	$\frac{10.7}{16}$ = 0.669																	
			Mole ratio (divide through by smallest value)	$\frac{6.667}{0.669}$ = 9.96	$\frac{9.300}{0.669}$ 13.90	$\frac{0.669}{0.669}$ 1.00																	
Round to Whole Number	10	14	1																				
25	C	58	<p>1st adjacent hydrogen</p> <p>2nd adjacent hydrogen</p> <p>3rd adjacent hydrogen</p> <p>This is not an adjacent hydrogen as it is attached to same carbon</p>																				
			<p>singlet no adjacent hydrogens</p> <p>Doublet 1 adjacent hydrogen</p> <p>Triplet 2 adjacent hydrogens</p> <p>quartet 3 adjacent hydrogens</p> <p>quintet 4 adjacent hydrogens</p>																				
26	A	66	<input checked="" type="checkbox"/> A both are agonists as both bind with a receptor and stimulate a response <input checked="" type="checkbox"/> B Buprenorphine is an agonist as it stimulates receptors and produces a response <input checked="" type="checkbox"/> C Pramipexole is an agonist as it stimulates the nerves cells to make a response <input checked="" type="checkbox"/> D Pramipexole is an agonist as it stimulates the nerves cells to make a response																				
27	D	80	1 minute = 6litres air \therefore 10minutes = 60litres air 0.03ppm = 0.03mg per 1litre air 1 litre of air \longleftrightarrow 0.03mg hydrogen sulphide 60 litres of air \longleftrightarrow 0.03mg hydrogen sulphide $\times \frac{60}{1}$ = 1.8mg																				
28	C	87	<input checked="" type="checkbox"/> A Thin layer chromatography separates chemicals with different polarities/size <input checked="" type="checkbox"/> B Heating under reflux allows volatile chemicals to react without reactant escape <input checked="" type="checkbox"/> C Recrystallisation allows chemical to be extracted from impurities due to solubility in a second solvent <input checked="" type="checkbox"/> D Distillation separates chemicals with different boiling points																				
29	D	70	<input checked="" type="checkbox"/> A water is less dense so should be the top layer <input checked="" type="checkbox"/> B water is less dense so should be the top layer <input checked="" type="checkbox"/> C $K = \frac{2}{8} = 0.25$ <input checked="" type="checkbox"/> D $K = \frac{8}{2} = 4$																				
30	A	52	<input checked="" type="checkbox"/> A Complexometric analysis is used to calculate the calcium ions in milk <input checked="" type="checkbox"/> B Gravimetric analysis is used to calculate the chloride ions in sea water <input checked="" type="checkbox"/> C Volumetric analysis is used to calculate ethanoic acid in vinegar <input checked="" type="checkbox"/> D Volumetric analysis is used to calculate ethanol in wine																				
















2019 Adv Higher Chemistry Marking Scheme

Long Qu	Answer	Reasoning						
1a	Any <u>one</u> of the 2p electrons circled	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%; text-align: center;">n=2 Electron in Shell 2</td> <td style="width: 25%; text-align: center;">l=1 Electron in p subshell</td> <td style="width: 25%; text-align: center;">m_l = -1 Electron in orbital -1 <small>(could be any of 3 orbitals in diagram)</small></td> <td style="width: 25%; text-align: center;">m_s = +$\frac{1}{2}$ Electron has +$\frac{1}{2}$ spin <small>(could be either up or down in diagram)</small></td> </tr> </table>	n=2 Electron in Shell 2	l=1 Electron in p subshell	m _l = -1 Electron in orbital -1 <small>(could be any of 3 orbitals in diagram)</small>	m _s = + $\frac{1}{2}$ Electron has + $\frac{1}{2}$ spin <small>(could be either up or down in diagram)</small>		
n=2 Electron in Shell 2	l=1 Electron in p subshell	m _l = -1 Electron in orbital -1 <small>(could be any of 3 orbitals in diagram)</small>	m _s = + $\frac{1}{2}$ Electron has + $\frac{1}{2}$ spin <small>(could be either up or down in diagram)</small>					
1b	103	$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $= 127 - 298 \times 79.4/1000$ $= 127 - 23.7$ $= 103.3 \text{ kJ mol}^{-1}$						
1c	3.20×10^{-5}	$\Delta G^\circ = -2.30 \times R \times T \times \log_{10}K$ $\log_{10}K = \frac{\Delta G^\circ}{-2.30 \times R \times T}$ $\log_{10}K = \frac{25.6}{-2.30 \times 8.31 \times 10^{-3} \times 298}$ $\log_{10}K = -4.49$ $K = 10^{-4.49} = 3.20 \times 10^{-5}$						
2a	Number of moles of reactant involved in the rate determining step	Rate determining step is the slowest step in reaction mechanism. The number of moles of each reactant in rate determining step decides the order for each reactant <table border="1" style="width: 100%; border-collapse: collapse; margin-top: 5px;"> <tr> <td style="width: 33%; text-align: center;">0 moles of reactant in RDS Zero Order</td> <td style="width: 33%; text-align: center;">1 mole of reactant in RDS 1st Order</td> <td style="width: 33%; text-align: center;">2 moles of reactant in RDS 2nd Order</td> </tr> </table>	0 moles of reactant in RDS Zero Order	1 mole of reactant in RDS 1 st Order	2 moles of reactant in RDS 2 nd Order			
0 moles of reactant in RDS Zero Order	1 mole of reactant in RDS 1 st Order	2 moles of reactant in RDS 2 nd Order						
2b(i)	2 nd order	1 mol of H ₂ O ₂ in rate determining step ∴ order [H ₂ O ₂] ¹ 1 mol of I ⁻ in rate determining step ∴ order [I ⁻] ¹ Overall order = 1 + 1 = 2						
2b(ii)	Rate = k [H ₂ O ₂] [I ⁻]	Rate = k [H ₂ O ₂] ¹ × [I ⁻] ¹ = k [H ₂ O ₂] [I ⁻]						
2c	$\text{H}_2\text{O}_2 + 2\text{I}^- + 2\text{H}_3\text{O}^+ \rightarrow \text{I}_2 + 4\text{H}_2\text{O}$	Step 1 H ₂ O ₂ + I ⁻ → IO⁻ + H ₂ O Step 2 IO⁻ + H ₃ O ⁺ → HOI + H ₂ O Step 3 HOI + H ₃ O ⁺ + I ⁻ → I ₂ + 2H ₂ O Overall H ₂ O ₂ + 2I ⁻ + 2H ₃ O ⁺ → I ₂ + 4H ₂ O						
3a	Answer to include:	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 10%;">1st Mark</td> <td style="width: 40%;">5cm³</td> <td style="width: 50%;">As a 1/10 dilution needs to take place, 5cm³ is transferred to the 50cm³ flask</td> </tr> <tr> <td>2nd Mark</td> <td>Correct use of pipette Making Up to line with deionised water</td> <td>An accurate method of measuring 5cm³ need to be used e.g. pipette The volumetric/standard flask must be filled to the line for accuracy</td> </tr> </table>	1 st Mark	5cm ³	As a 1/10 dilution needs to take place, 5cm ³ is transferred to the 50cm ³ flask	2 nd Mark	Correct use of pipette Making Up to line with deionised water	An accurate method of measuring 5cm ³ need to be used e.g. pipette The volumetric/standard flask must be filled to the line for accuracy
1 st Mark	5cm ³	As a 1/10 dilution needs to take place, 5cm ³ is transferred to the 50cm ³ flask						
2 nd Mark	Correct use of pipette Making Up to line with deionised water	An accurate method of measuring 5cm ³ need to be used e.g. pipette The volumetric/standard flask must be filled to the line for accuracy						
3b(i)	Deionised water	The control experiment has a cuvette containing deionised water and the absorbance of this is the setting for zero absorbance.						
3b(ii)	Unknown's Absorbance must not be outwith calibration range	The calibration curve is only applicable to concentrations between the highest and lowest concentrations used in the calibration curve. Diluting a sample can bring a more concentrated unknown into the usable range of the calibration curve. It is unknown what happens to concentrations beyond the calibration range with any certainty.						
3b(iii)	71%	Absorbance (diluted sample) = 0.34 ∴ concentration of Cu ²⁺ = 0.032 mol l ⁻¹ ∴ concentration of Cu ²⁺ in original sample = 0.064 mol l ⁻¹ no. of mol in 250cm ³ = volume × concentration = 0.25litres × 0.064 mol l ⁻¹ = 0.016mol mass = no. of mol × gfm = 0.016 × 63.5 = 1.016g $\% \text{ mass} = \frac{\text{mass of Cu}}{\text{mass of screw}} \times 100 = \frac{1.016}{1.43} \times 100 = 71.0\%$						

4a(i)	Proton or H ⁺ acceptor	Acid	Donates H ⁺																		
		Base	Accepts H ⁺																		
		Conjugate Acid	Formed when Base accepts H ⁺																		
		Conjugate Base	Formed when Acid loses H																		
4a(ii)	<table border="1"> <tr><td>Acid</td><td>H₂O₂</td></tr> <tr><td>Conjugate Base</td><td>HO₂⁻</td></tr> </table>	Acid	H ₂ O ₂	Conjugate Base	HO ₂ ⁻	$\text{H}_2\text{O}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{HO}_2^-$ <p>Acid Base Conjugate Acid Conjugate Base Donates H⁺ Accepts H⁺ Formed when Base accepts H⁺ Formed when Acid loses H⁺</p>	or														
	Acid	H ₂ O ₂																			
Conjugate Base	HO ₂ ⁻																				
<table border="1"> <tr><td>Acid</td><td>H₃O⁺</td></tr> <tr><td>Conjugate Base</td><td>H₂O</td></tr> </table>	Acid	H ₃ O ⁺	Conjugate Base	H ₂ O	$\text{H}_3\text{O}^+ + \text{HO}_2^- \rightleftharpoons \text{H}_2\text{O}_2 + \text{H}_2\text{O}$ <p>Acid Base Conjugate Acid Conjugate Base Donates H⁺ Accepts H⁺ Formed when Base accepts H⁺ Formed when Acid loses H⁺</p>																
Acid	H ₃ O ⁺																				
Conjugate Base	H ₂ O																				
4b	One answer from:	B(OH) ₃ accepts a pair of non-bonding electrons and water donates a pair of non-bonding electrons	B(OH) ₃ accepts a pair of non-bonding electrons from water.	Water donates a pair of non-bonding electrons to B(OH) ₃																	
4c(i)	Increasing number of chlorine increases strength of acid	Increasing the number of chlorines increases the value of K _a . The higher the value of K _a the stronger the acid (NB The higher the value of pK _a the weaker the acid)																			
4c(ii) Part A	0.08	$\text{gfm CH}_2\text{COOH} = (2 \times 12) + (3 \times 1) + (2 \times 16) + (1 \times 35.5) = 24 + 3 + 32 + 35.5 = 94.5\text{g}$ $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{1.89}{94.5} = 0.02\text{mol}$ $\text{concentration} = \frac{\text{no. of mol}}{\text{volume}} = \frac{0.02 \text{ mol}}{0.25 \text{ litres}} = 0.08 \text{ mol l}^{-1}$																			
4c(ii) Part B	2.0	$\text{pH} = \frac{1}{2} \text{pK}_a - \frac{1}{2} \log c$ $\text{pH} = \frac{1}{2} \times -\log_{10} K_a - \frac{1}{2} \log_{10} c$ $\text{pH} = -\frac{1}{2} \times \log_{10} (1.3 \times 10^{-3}) - \frac{1}{2} \log_{10} (0.08)$ $\text{pH} = -\frac{1}{2} \times (-2.89) - \frac{1}{2} \times (-1.10)$ $\text{pH} = 1.44 - (-0.55)$ $\text{pH} = 1.99$																			
4d	Open Question to include:	3 mark answer	2 mark answer	1 mark answer																	
		Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.																	
5a	Heat or high temperature	Electrons become excited in sodium by absorbing heat energy which promotes an electron to a higher energy level. Light of specific wavelength is emitted corresponding to the exact energy difference between the upper energy level the electron drops down from to the lower energy level the electron goes in to.																			
5b(i)	Hexaaquazinc(II)	Hexaaquazinc(II) = [Zn(H₂O)₆]²⁺																			
		<table border="1"> <tr> <td>no. of ligands</td> <td>H₂O ligand</td> <td>metal name</td> <td>Charge on metal ion</td> </tr> </table>	no. of ligands	H ₂ O ligand	metal name	Charge on metal ion	Neutral ligands include:	Negative Ligands include:	Central Ion:	Charge:											
no. of ligands	H ₂ O ligand	metal name	Charge on metal ion																		
		<table border="1"> <tr><td>Ligand</td><td>Name</td></tr> <tr><td>OH₂</td><td>aqua</td></tr> <tr><td>NH₃</td><td>ammine</td></tr> <tr><td>CO</td><td>carbonyl</td></tr> </table>	Ligand	Name	OH ₂	aqua	NH ₃	ammine	CO	carbonyl	<table border="1"> <tr><td>Ligand</td><td>Name</td></tr> <tr><td>Chloride Cl⁻</td><td>chlorido</td></tr> <tr><td>Cyanide CN⁻</td><td>cyanido</td></tr> <tr><td>Nitrite NO₂⁻</td><td>nitrito</td></tr> </table>	Ligand	Name	Chloride Cl ⁻	chlorido	Cyanide CN ⁻	cyanido	Nitrite NO ₂ ⁻	nitrito	Positive Complex: metals keep their name Negative Complex: Metals end in ATE e.g. Cuprate, Ferrate, Cobaltate	Charge of central ion is converted into roman numerals and put in brackets
Ligand	Name																				
OH ₂	aqua																				
NH ₃	ammine																				
CO	carbonyl																				
Ligand	Name																				
Chloride Cl ⁻	chlorido																				
Cyanide CN ⁻	cyanido																				
Nitrite NO ₂ ⁻	nitrito																				
5b(ii)	Answer to include:	1 st Mark		2 nd Mark																	
		Full or complete d subshell or 10 d electrons or 3d ¹⁰ or no d-d transitions possible		It does not absorb visible light or It only absorbs ultraviolet/UV light																	

5c(i)	$2.04 \times 10^{-16} \text{ J}$	$E = h \times f$ $E = 6.63 \times 10^{-34} \text{ J s} \times 3.08 \times 10^{17} \text{ s}^{-1}$ $E = 2.04 \times 10^{-16} \text{ J}$ <p>NB: The question does not require you to multiply by L (6.02×10^{23}) as the units required are J not J mol^{-1} or kJ mol^{-1}</p>																				
5c(ii)	19	$1 \text{ J} \longleftrightarrow 6.24 \times 10^{18} \text{ eV}$ $2.04 \times 10^{-16} \text{ J} \longleftrightarrow 6.24 \times 10^{18} \text{ eV} \times 2.04 \times 10^{-16} / 1$ $= 1273 \text{ eV}$ <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="text-align: right;">Binding Energy</td> <td style="text-align: center;">=</td> <td style="text-align: center;">Energy of electromagnetic radiation</td> <td style="text-align: center;">-</td> <td style="text-align: center;">Kinetic Energy of electron emitted</td> </tr> <tr> <td style="text-align: right;">E_b</td> <td style="text-align: center;">=</td> <td style="text-align: center;">E</td> <td style="text-align: center;">-</td> <td style="text-align: center;">E_k</td> </tr> <tr> <td style="text-align: right;">E_b</td> <td style="text-align: center;">=</td> <td style="text-align: center;">1273 eV</td> <td style="text-align: center;">-</td> <td style="text-align: center;">1254 eV</td> </tr> <tr> <td style="text-align: right;">E_b</td> <td style="text-align: center;">=</td> <td style="text-align: center;">19 eV</td> <td></td> <td></td> </tr> </table>	Binding Energy	=	Energy of electromagnetic radiation	-	Kinetic Energy of electron emitted	E_b	=	E	-	E_k	E_b	=	1273 eV	-	1254 eV	E_b	=	19 eV		
Binding Energy	=	Energy of electromagnetic radiation	-	Kinetic Energy of electron emitted																		
E_b	=	E	-	E_k																		
E_b	=	1273 eV	-	1254 eV																		
E_b	=	19 eV																				
6a	6.4	<p>Total Dichromate in flask: no. of mol = volume \times concentration = $0.025 \text{ litres} \times 0.010 \text{ mol l}^{-1} = 2.5 \times 10^{-4} \text{ mol}$</p> <p>Dichromate left at end of reaction: no. of mol = $1.65 \times 10^{-4} \text{ mol}$ (in question)</p> <p>Dichromate which reacted with ethanol: no. of mol = $2.5 \times 10^{-4} \text{ mol} - 1.65 \times 10^{-4} \text{ mol} = 8.5 \times 10^{-5} \text{ mol}$</p> $3\text{C}_2\text{H}_5\text{OH} + 2\text{Cr}_2\text{O}_7^{2-} + 16\text{H}^+ \longrightarrow 3\text{CH}_3\text{COOH} + 4\text{Cr}^{3+} + 11\text{H}_2\text{O}$ <table style="width: 100%; border-collapse: collapse; margin-left: 20px;"> <tr> <td style="text-align: center;">3mol</td> <td style="text-align: center;">2mol</td> <td></td> <td></td> <td></td> </tr> <tr> <td style="text-align: center;">$1.28 \times 10^{-4} \text{ mol}$</td> <td style="text-align: center;">$8.5 \times 10^{-5} \text{ mol}$</td> <td></td> <td></td> <td></td> </tr> </table> $1 \text{ cm}^3 \text{ diluted vodka} \longleftrightarrow 1.28 \times 10^{-4} \text{ mol}$ $1000 \text{ cm}^3 \text{ diluted vodka} \longleftrightarrow 1.28 \times 10^{-4} \text{ mol} \times 1000 / 1$ $= 0.128 \text{ mol}$ <p>As 1000 cm^3 diluted vodka was made from 20.0 cm^3 of vodka $\therefore 20 \text{ cm}^3$ undiluted vodka $\longleftrightarrow 0.128 \text{ mol}$</p> $\text{concentration} = \frac{\text{no. of mol}}{\text{volume}} = \frac{0.128 \text{ mol}}{0.020 \text{ litres}} = 6.4 \text{ mol l}^{-1}$	3mol	2mol				$1.28 \times 10^{-4} \text{ mol}$	$8.5 \times 10^{-5} \text{ mol}$													
3mol	2mol																					
$1.28 \times 10^{-4} \text{ mol}$	$8.5 \times 10^{-5} \text{ mol}$																					
6b	To ensure all ethanol is reacted	This experiment is a back titration. An excess of acidified potassium dichromate is added to ethanol solution and all the ethanol will react with acidified potassium dichromate solution and the left over acidified potassium dichromate can be determined by volumetric analysis.																				
6c	Impurities in vodka reacting with dichromate	Other chemicals in the vodka may be able to react with acidified dichromate (a powerful oxidising agent).																				
6d	Use standard solution of ethanol	A standard solution of ethanol could be used as a control to measure the experimentally-determined value of this ethanol solution against the known value of the standard solution.																				
7a	Under suction/vacuum	A Buchner funnel and suction pump is used to draw the filtrate through the filter paper in the Buchner funnel quicker than gravity alone.																				
7b(i)	End on overlap of two atomic orbitals	A sigma bond is formed by the end on overlap of two orbitals. A pi bond is formed by side overlap of two unhybridized p orbitals																				
7b(ii)	Mixing an s atomic orbital with two p atomic orbitals	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <thead> <tr> <th style="padding: 5px;">Type of Hybridisation</th> <th style="padding: 5px;">sp^3</th> <th style="padding: 5px;">sp^2</th> <th style="padding: 5px;">sp</th> </tr> </thead> <tbody> <tr> <td style="padding: 5px;">Description</td> <td style="padding: 5px;">One s and three p orbitals mix</td> <td style="padding: 5px;">One s and two p orbitals mix</td> <td style="padding: 5px;">One s and one p orbitals mix</td> </tr> </tbody> </table>	Type of Hybridisation	sp^3	sp^2	sp	Description	One s and three p orbitals mix	One s and two p orbitals mix	One s and one p orbitals mix												
Type of Hybridisation	sp^3	sp^2	sp																			
Description	One s and three p orbitals mix	One s and two p orbitals mix	One s and one p orbitals mix																			

7c	Answer to include:	1 st Mark		2 nd Mark			
		Electrons promoted/move from HOMO to LUMO		Blue/green light is absorbed or Complementary colour to red absorbed			
7d(i)	One answer from:	Similar polarities or both polar or similar intermolecular forces	Forms hydrogen bonds with alizarin	Does not react with alizarin	Volatile or low boiling point or evaporates easily	Dissolves alizarin but not other substances (in the root)	
7d(ii) Part A	Answer to include:	1 st Mark		2 nd Mark			
		IR makes bonds vibrate/bend/stretch		Different bonds/functional groups absorb at different wavenumber/wavelength/frequency/energy			
7d(ii) Part B	One or both -OH groups circled	From page 14 of data booklet:					
		Wave Number Range/cm ⁻¹ 3570-3200	Type of Compound Alcohols and Phenols	Infra-red Absorption due to Hydrogen bonded O - H stretch			
The -OH hydroxyl group can be described as phenols as they are attached to a benzene ring.							
7d(ii) Part C (I)	2.96x10 ⁻⁶ m	Wavelength = $\frac{1}{\text{Wavenumber}} = \frac{1}{3395\text{cm}^{-1}} = 2.96 \times 10^{-4} \text{ cm} = 2.96 \times 10^{-6} \text{ m}$					
7d(ii) Part C (II)	40.65kJ mol ⁻¹	$E = \frac{L \times h \times c}{\lambda} = \frac{6.02 \times 10^{23} \text{ mol}^{-1} \times 6.63 \times 10^{-34} \text{ J s} \times 3 \times 10^8 \text{ m s}^{-1}}{2.96 \times 10^{-6} \text{ m}}$ $= 40651 \text{ J mol}^{-1}$ $= 40.65 \text{ kJ mol}^{-1}$					
8	Open Question to include:	3 mark answer			2 mark answer		1 mark answer
		Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.			Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.		Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
9a(i)	Either circle on diagram: 	A chiral carbon has four different chemical groups attached to a central carbon. Both circled carbons have four bond but due to being skeletal formula diagrams the 4th bond (a C-H bond) being the 4 th different group attached to the central carbon is not drawn as it is a C-H bond.					
9a(ii)	Non-superimposable mirror images	Non-superimposable Mirror Images Images shown are direct mirror images of each other and are therefore are optical isomers (enantiomers)			Non-superimposable Mirror Images The following are optical isomers as Groups W and X are in the same position but Groups Y and Z are in different position.		
							
9b(i)	(nucleophilic) substitution	The lone pair on the pair of the N in NH ₃ is nucleophilically attracted to the δ ⁺ on the C of the polar C-Br bond. The NH ₃ is added onto the carbon as the Br atom leaves as a Br ⁻ ion. An H in the -NH ₃ ⁺ group then leaves as an H ⁺ ion leaving a -NH ₂ amine group attached to the carbon the Br previously was attached to.					

9b(ii)	7.58g	<p>1-phenylpropanone \longrightarrow [intermediate] \longrightarrow cathinone</p> <p>1mol 1mol 134g 149g 9.50g $149g \times 9.50/134$ $= 10.56g$</p> <p>$\% \text{Yield} = \frac{\text{Actual}}{\text{Theoretical}} \times 100 \therefore \text{Actual} = \frac{\% \text{ Yield} \times \text{Theoretical}}{100} = \frac{71.8 \times 10.56}{100} = 7.58g$</p>				
10a	Ether	Ethers have a carbon-oxygen-carbon group as their functional group. Be careful not to confuse a C-O-C group with an ester group which also has a carbonyl C=O group in one end.				
10b(i)	One from:					
10b(ii)	2-methoxy-2-methylpropane					
10c(i)	methanol	<p>Group 1 metals react with alcohols to produce alkoxides and hydrogen:</p> <p>sodium + methanol \longrightarrow sodium methoxide + hydrogen $2\text{Na} + 2\text{CH}_3\text{OH} \longrightarrow 2\text{Na}^+\text{CH}_3\text{O}^- + \text{H}_2$</p>				
10c(ii)	<p><u>1st Mark</u> Correct Carbocation intermediate</p> <p><u>2nd Mark</u> Correct use of <u>both</u> curly arrows</p>					
10c(iii)	One answer from:	<table border="1" style="width:100%"> <tr> <td style="width:50%">Formation of stable (tertiary) carbocation</td> <td style="width:50%">Steric hindrance of three methyl -CH₃ groups prevents nucleophilic attack</td> </tr> </table>	Formation of stable (tertiary) carbocation	Steric hindrance of three methyl -CH ₃ groups prevents nucleophilic attack		
Formation of stable (tertiary) carbocation	Steric hindrance of three methyl -CH ₃ groups prevents nucleophilic attack					
10d	Any one from:	<table border="1" style="width:100%"> <tr> <td style="text-align:center">  pentan-2-ol </td> <td style="text-align:center">  3-methylbutan-2-ol </td> <td style="text-align:center">  2-methylbutan-1-ol </td> <td style="text-align:center">  2-methoxybutane </td> </tr> </table>	 pentan-2-ol	 3-methylbutan-2-ol	 2-methylbutan-1-ol	 2-methoxybutane
 pentan-2-ol	 3-methylbutan-2-ol	 2-methylbutan-1-ol	 2-methoxybutane			
10e	One line at 1.5-0.9 with relative intensity 9	<p>The peak drawn is 3.6-3.7 and a relative intensity of 3 \therefore peak caused -O-CH₃ group (3xH = relative intensity 3)</p> <p>Other Peak must contain tertiary -C(CH₃) group \therefore Line at chemical shift 1.5-0.9 for R₃C- from table \therefore Relative intensity of 9 as there are 9 H atoms in group.</p>				