



JABchem



Not to be shared without the copyright holder's permission

Past Papers Higher Chemistry

2018 Marking Scheme

Grade Awarded	Mark Required		% candidates achieving grade
	(/120)	%	
A	80+	66.6%	28.3%
B	67+	55.8%	25.3%
C	54+	45.0%	23.0%
D	47+	39.2%	9.5%
No award	<47	<39.2%	13.9%




Section:	Multiple Choice	Extended Answer	Assignment
Average Mark:	12.1 /20	42.9 /80	13.1 /20

2018 Higher Chemistry Marking Scheme

MC Qu	Answer	% Pupils Correct	Reasoning								
1	B		<input checked="" type="checkbox"/> A Forward reaction has an activation energy of +40kJ mol ⁻¹ but always positive <input checked="" type="checkbox"/> B Enthalpy change is difference between P and R and downhill means exothermic <input checked="" type="checkbox"/> C This is enthalpy change for forward reaction from R to P (uphill = endothermic) <input checked="" type="checkbox"/> D This is the activation energy for the reverse reaction (P to top of hill)								
2	A		$\text{Rate} = \frac{1}{t} \therefore t = \frac{1}{\text{rate}} = \frac{1}{100 \text{ s}} = 0.01 \text{ s}^{-1}$								
3	D		<input checked="" type="checkbox"/> A Decrease in temp decreases number of collisions with energy greater than E _a <input checked="" type="checkbox"/> B Activation Energy (E _a) is independent of temperature <input checked="" type="checkbox"/> C Activation Energy (E _a) is independent of temperature <input checked="" type="checkbox"/> D No change to activation energy & number of collisions decrease as temp decreases								
4	C		$\begin{array}{l} \textcircled{1} \quad \text{Al}^+(\text{g}) \rightarrow \text{Al}^{2+}(\text{g}) + \text{e}^- \quad \Delta H=1817\text{kJ} \\ \textcircled{2} \quad \text{Al}^{2+}(\text{g}) \rightarrow \text{Al}^{3+}(\text{g}) + \text{e}^- \quad \Delta H=2745\text{kJ} \\ \text{Add } \textcircled{1}+\textcircled{2} \quad \begin{array}{l} \text{Al}^+(\text{g}) \rightarrow \cancel{\text{Al}^{2+}(\text{g})} + \text{e}^- \\ \cancel{\text{Al}^{2+}(\text{g})} \rightarrow \text{Al}^{3+}(\text{g}) + \text{e}^- \\ \hline \text{Al}^+(\text{g}) \rightarrow \text{Al}^{3+}(\text{g}) + 2\text{e}^- \quad \Delta H=4562\text{kJ} \end{array} \end{array}$								
5	D		<input checked="" type="checkbox"/> A Boron forms covalent network. No molecules to have London Dispersion Forces between <input checked="" type="checkbox"/> B Neon is a noble gas and contains no covalent bonds <input checked="" type="checkbox"/> C Sodium is an alkali metal and contains metallic bonds between atoms <input checked="" type="checkbox"/> D Sulphur has covalent bonds within its S ₈ rings and LDF between molecules.								
6	C		<table border="1" style="width: 100%; text-align: center;"> <tr> <td>—O—H</td> <td>$\begin{array}{c} \text{O} \\ \\ \text{—C—OH} \end{array}$</td> <td>$\begin{array}{c} \text{O} \\ \\ \text{—C—O—} \end{array}$</td> <td>$\begin{array}{c} \text{O} \\ \\ \text{—C—} \end{array}$</td> </tr> <tr> <td>hydroxyl group</td> <td>carboxyl group</td> <td>ester group</td> <td>carbonyl group</td> </tr> </table>	—O—H	$\begin{array}{c} \text{O} \\ \\ \text{—C—OH} \end{array}$	$\begin{array}{c} \text{O} \\ \\ \text{—C—O—} \end{array}$	$\begin{array}{c} \text{O} \\ \\ \text{—C—} \end{array}$	hydroxyl group	carboxyl group	ester group	carbonyl group
—O—H	$\begin{array}{c} \text{O} \\ \\ \text{—C—OH} \end{array}$	$\begin{array}{c} \text{O} \\ \\ \text{—C—O—} \end{array}$	$\begin{array}{c} \text{O} \\ \\ \text{—C—} \end{array}$								
hydroxyl group	carboxyl group	ester group	carbonyl group								
7	A		<input checked="" type="checkbox"/> A main chain has 4 carbons, carboxyl functional group and methyls on C ₂ , C ₂ and C ₃ <input checked="" type="checkbox"/> B side groups incorrectly numbered as C ₂ has two methyl groups attached. <input checked="" type="checkbox"/> C carbon side groups e.g. methyl groups never appear on carbon number 1 <input checked="" type="checkbox"/> D longest chain with functional group is four carbons ∴ ends in butanoic acid								
8	D		<input checked="" type="checkbox"/> A CH ₃ H ₂ CH(OH)CH ₂ CH ₃ is pentan-3-ol so cannot be an isomer of pentan-3-ol <input checked="" type="checkbox"/> B Structure has formula C ₆ H ₁₃ OH so has different formula and not an isomer <input checked="" type="checkbox"/> C Structure has formula C ₅ H ₉ OH so has different formula and not an isomer <input checked="" type="checkbox"/> D Structure is pentan-1-ol, has same formula and a different structure								
9	A		4-methylpentan-2-ol C ₆ H ₁₄ O → 4-methylpentan-2-one C ₆ H ₁₂ O ∴ loss of 2xH atoms <input checked="" type="checkbox"/> A Loss of 2g per mole would be losing 2xH atoms from 1mol of C ₆ H ₁₄ O <input checked="" type="checkbox"/> B Gain of 2g per mole would be gaining 2xH atoms on 1mol of C ₆ H ₁₄ O <input checked="" type="checkbox"/> C Loss of 16g per mole would be losing 1xO atom from 1mol of C ₆ H ₁₄ O <input checked="" type="checkbox"/> D Gain of 16g per mole would be gaining 1xO atom on 1mol of C ₆ H ₁₄ O								
10	B		<input checked="" type="checkbox"/> A All amino acids are necessary for building the different types of proteins <input checked="" type="checkbox"/> B Essential amino acids are the amino acids humans must acquire in their diet. <input checked="" type="checkbox"/> C Plants produce the essential amino acids that humans consume in their diet <input checked="" type="checkbox"/> D All types of amino acids are produced when a protein is hydrolysed								
11	D		$\begin{array}{l} n \text{ CO} + (2n+1) \text{ H}_2 \rightarrow n \text{ H}_2\text{O} + \text{C}_n\text{H}_x \\ n \text{ CO} + 2n \text{ H}_2 + \text{H}_2 \rightarrow n \text{ H}_2\text{O} + \text{C}_n\text{H}_x \\ n \text{ CO} + n \text{ H}_2 + n \text{ H}_2 + \text{H}_2 \rightarrow n \text{ H}_2\text{O} + \text{C}_n\text{H}_x \\ n \text{ C} + n \text{ H}_2 + n \text{ H}_2 + \text{H}_2 \rightarrow n \text{ H}_2 + \text{C}_n\text{H}_x \\ n \text{ C} + 2n \text{ H} + 2\text{H} \rightarrow \quad \quad \quad + \text{C}_n\text{H}_x \\ n \text{ C} + 2n+2 \text{ H} \rightarrow \quad \quad \quad + \text{C}_n\text{H}_x \end{array}$ <p>n Carbon atoms has 2n+2 Hydrogen atoms available: general formula = C_nH_{2n+2}</p>								

12	D	<p>1mol Na⁺Cl⁻ contains 1mol Cl⁻ ions ∴ 0.6mol Na⁺Cl⁻ contains 0.6mol Cl⁻ ions</p> <p>1mol Na⁺Cl⁻ contains 1mol Na⁺ ions ∴ 0.6mol Na⁺Cl⁻ contains 0.6mol Na⁺ ions</p>	<p>1mol (Na⁺)₂SO₄²⁻ contains 1mol SO₄²⁻ ions ∴ 0.2mol (Na⁺)₂SO₄²⁻ contains 0.2mol SO₄²⁻ ions</p> <p>1mol (Na⁺)₂SO₄²⁻ contains 2mol Na⁺ ions ∴ 0.2mol (Na⁺)₂SO₄²⁻ contains 0.4mol Na⁺ ions</p>
13	A	<p><input checked="" type="checkbox"/> A 0.20g of H₂ ∴ no. of mol = $\frac{\text{mass}}{\text{gfm}} = \frac{0.20}{2} = 0.10\text{mol}$ ∴ largest volume</p> <p><input checked="" type="checkbox"/> B 0.44g of CO₂ ∴ no. of mol = $\frac{\text{mass}}{\text{gfm}} = \frac{0.44}{44} = 0.01\text{mol}$ ∴ smallest volume</p> <p><input checked="" type="checkbox"/> C 0.60g of Ne ∴ no. of mol = $\frac{\text{mass}}{\text{gfm}} = \frac{0.60}{20} = 0.03\text{mol}$</p> <p><input checked="" type="checkbox"/> D 0.80g of Ar ∴ no. of mol = $\frac{\text{mass}}{\text{gfm}} = \frac{0.80}{2} = 0.02\text{mol}$</p>	
14	A	$3\text{CuO}_{(s)} + 2\text{NH}_{3(g)} \longrightarrow 2\text{Cu}_{(s)} + \text{N}_{2(g)} + 3\text{H}_2\text{O}_{(l)}$ <p style="text-align: center;"> 3mol 2mol 2mol 1mol 3mol negligible volume 2vol negligible volume 1vol negligible volume - 100cm³ - 50cm³ - </p>	
15	D	<p><input checked="" type="checkbox"/> A Hydrogen gas (H₂) would have no effect as it is neither a reactant nor product</p> <p><input checked="" type="checkbox"/> B HCl(g) would dissolve in water to form acid sending equilibrium to left to remove H⁺ ions</p> <p><input checked="" type="checkbox"/> C Cl⁻ ions added so equilibrium would move to left to remove extra Cl⁻ ions</p> <p><input checked="" type="checkbox"/> D OH⁻ ions would neutralise H⁺ ions sending equilibrium to right to replace H⁺ ions.</p>	
16	C	<p>gfm C₄H₉OH = 72 mass = 3.6g</p> <p style="text-align: right;">no. of mol = $\frac{\text{mass}}{\text{gfm}} = \frac{3.6}{72} = 0.05\text{mol}$</p> <p>0.05mol C₄H₉OH releases 124kJ 1mol C₄H₉OH releases 124kJ × 1/0.05 = -2480kJ mol⁻¹</p>	
17	B	<p>① C(s) + O₂(g) → CO₂ ΔH=-394kJ</p> <p>② CO(g) + $\frac{1}{2}$O₂(g) → CO₂(g) ΔH=-283kJ</p> <p>① C(s) + O₂(g) → CO₂ ΔH=-394kJ</p> <p>② × -1 CO₂(g) → CO(g) + $\frac{1}{2}$O₂(g) ΔH=+283kJ</p> <p>Add ①+②' C(s) + $\frac{1}{2}$O₂(g) → CO₂ CO₂(g) → CO(g) + $\frac{1}{2}$O₂(g)</p> <p>C(s) + $\frac{1}{2}$O₂(g) → CO(g) ΔH=-111kJ</p>	
18	C	<p><input checked="" type="checkbox"/> A Cr³⁺ + 3e⁻ → Cr is higher in the electrochemical series than the equation SO₄²⁻ + 2H⁺ + 2e⁻ → SO₃²⁻ + H₂O is too low to turn Cr³⁺ into Cr</p> <p><input checked="" type="checkbox"/> B Al³⁺ + 3e⁻ → Al is higher in the electrochemical series than the equation SO₄²⁻ + 2H⁺ + 2e⁻ → SO₃²⁻ + H₂O is too low to turn Al³⁺ into Al.</p> <p><input checked="" type="checkbox"/> C Fe³⁺ + e⁻ → Fe²⁺: Fe³⁺ ions will reduce to Fe²⁺ ions and is lower in electrochemical series than SO₄²⁻ + H₂O → SO₄²⁻ + 2H⁺ + 2e⁻ (reversed as it is higher in ECS)</p> <p><input checked="" type="checkbox"/> D Sn⁴⁺ + 2e⁻ → Sn²⁺ is higher in electrochemical series than the equation SO₄²⁻ + 2H⁺ + 2e⁻ → SO₃²⁻ + H₂O is too low to turn Sn⁴⁺ into Sn²⁺.</p>	
19	B	<p><u>Step 1:</u> Write down main species in reaction NO₃⁻ → NO</p> <p><u>Step 2:</u> Balance all atoms other than O or H NO₃⁻ → NO</p> <p><u>Step 3:</u> Balance O atoms by adding H₂O to the other side NO₃⁻ → NO + 2H₂O</p> <p><u>Step 4:</u> Balance H atoms by adding H⁺ to the other side NO₃⁻ + 4H⁺ → NO + 2H₂O</p> <p><u>Step 5:</u> Balance charge by adding electrons to the most positive side NO₃⁻ + 4H⁺ + 3e⁻ → NO + 2H₂O</p>	
20	B	<p>Increase in proportion of solid = Increase in rate of forward reaction</p> <p><input checked="" type="checkbox"/> A decrease in pressure increases the rate of the pressure-reducing reverse reaction</p> <p><input checked="" type="checkbox"/> B decrease in temp and increase in pressure both favour the forward reaction.</p> <p><input checked="" type="checkbox"/> C increase in temperature increases the rate of the endothermic reverse reaction</p> <p><input checked="" type="checkbox"/> D increase in temperature increases the rate of the endothermic reverse reaction</p>	

2018 Higher Chemistry Marking Scheme

Long Qu	Answer	Reasoning																
1a(i)	The attraction an atom/nucleus has for the electrons in a bond/shared electrons	Electronegativity is a measure of the attraction for the electrons in a bond by the nuclei at either end of that bond. Non-metals tend to have higher values of electronegativity and have a higher attraction for the electrons in a bond.																
1a(ii)	One answer from:	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; text-align: center;">Increased shielding/ more shielding (from additional shells)</td> <td style="width: 50%; text-align: center;">Covalent radius increases/atom size increases/more shells so attraction of the nucleus/protons for the (outer/shared) electrons decreases</td> </tr> </table>	Increased shielding/ more shielding (from additional shells)	Covalent radius increases/atom size increases/more shells so attraction of the nucleus/protons for the (outer/shared) electrons decreases														
Increased shielding/ more shielding (from additional shells)	Covalent radius increases/atom size increases/more shells so attraction of the nucleus/protons for the (outer/shared) electrons decreases																	
1b	Answer to include:	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; text-align: center;"><u>1 mark</u> (Intermolecular) forces/bonds increase (going down the group)</td> <td style="width: 33%; text-align: center;"><u>1 mark</u> London Dispersion Forces are the forces (broken) between the molecules</td> <td style="width: 33%; text-align: center;"><u>1 mark</u> The more electrons the stronger the London Dispersion Forces</td> </tr> </table>	<u>1 mark</u> (Intermolecular) forces/bonds increase (going down the group)	<u>1 mark</u> London Dispersion Forces are the forces (broken) between the molecules	<u>1 mark</u> The more electrons the stronger the London Dispersion Forces													
<u>1 mark</u> (Intermolecular) forces/bonds increase (going down the group)	<u>1 mark</u> London Dispersion Forces are the forces (broken) between the molecules	<u>1 mark</u> The more electrons the stronger the London Dispersion Forces																
2a	Increasing number of protons or increasing nuclear charge	Across a period the same outer shell is being filled by electrons. The number of protons in the nucleus is increasing as is the positive charge in the nucleus. The outer shell of electrons is pulled in closer to the nucleus due to electrostatic attraction making the atom size decrease.																
2b(i)		<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">Electronegativity</td> <td style="width: 25%;">Si=1.9</td> <td style="width: 25%;">P=2.2</td> <td style="width: 25%;">S=2.5</td> </tr> <tr> <td>Chlorine</td> <td>Cl=3.0</td> <td>Cl=3.0</td> <td>Cl=3.0</td> </tr> <tr> <td>Difference</td> <td>1.1</td> <td>0.8</td> <td>0.5</td> </tr> <tr> <td>Commentary</td> <td>Most polar bonds</td> <td>-</td> <td>Least polar bonds</td> </tr> </table>	Electronegativity	Si=1.9	P=2.2	S=2.5	Chlorine	Cl=3.0	Cl=3.0	Cl=3.0	Difference	1.1	0.8	0.5	Commentary	Most polar bonds	-	Least polar bonds
Electronegativity	Si=1.9	P=2.2	S=2.5															
Chlorine	Cl=3.0	Cl=3.0	Cl=3.0															
Difference	1.1	0.8	0.5															
Commentary	Most polar bonds	-	Least polar bonds															
2b(ii)	Answer to include:	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 15%; text-align: center;">1 mark</td> <td>Silicon tetrachloride and hexane are both non-polar</td> </tr> <tr> <td style="text-align: center;">1 mark</td> <td>Silicon tetrachloride is non-polar due to its tetrahedral shape (where polarities over molecule cancel out)</td> </tr> </table>	1 mark	Silicon tetrachloride and hexane are both non-polar	1 mark	Silicon tetrachloride is non-polar due to its tetrahedral shape (where polarities over molecule cancel out)												
1 mark	Silicon tetrachloride and hexane are both non-polar																	
1 mark	Silicon tetrachloride is non-polar due to its tetrahedral shape (where polarities over molecule cancel out)																	
2c(i)	Answer to include:	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 15%; text-align: center;">1 mark</td> <td>Silicon nitride is a covalent network</td> </tr> <tr> <td style="text-align: center;">1 mark</td> <td>Covalent bonds need to be broken before it will melt</td> </tr> </table>	1 mark	Silicon nitride is a covalent network	1 mark	Covalent bonds need to be broken before it will melt												
1 mark	Silicon nitride is a covalent network																	
1 mark	Covalent bonds need to be broken before it will melt																	
2c(ii)	17.9	$\text{atom economy} = \frac{140.3}{(3 \times 170.1) + (16 \times 17.0)} \times 100 = \frac{140.3}{510.3 + 272.0} \times 100 = 17.9\%$																
2d(i)	Answer to include:	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> <p><u>1 mark</u></p>  </div> <div style="text-align: center;"> <p><u>1 mark</u></p>  </div> </div>																
2d(ii)	To supply activation energy	Although the reaction is exothermic, sufficient energy to form the activated complex initially must be supplied for the activation energy barrier to be overcome. Once the reaction gets going the exothermic reaction will provide the heat energy to maintain the reaction.																
3a	water bath or heating mantle or hot plate	A flame-based method of heating should not be used as the reactants and products are volatile and flammable.																
3b	To condense any escaping gases	The reactants and products in the formation of esters can have low enough boiling points to evaporate and escape from the test tube. A cold surface like a small test tube filled with cold water will give a surface for the escaping gases to condense on. (Be careful, condensing gases give off heat which will rapidly heat up the water in the cold test tube)																
3c(i)	water	$\text{benzoic Acid} + \text{methanol} \rightarrow \text{methyl benzoate} + \text{water}$ $\text{C}_6\text{H}_5\text{COOH} + \text{CH}_3\text{OH} \rightarrow \text{C}_6\text{H}_5\text{COOCH}_3 + \text{H}_2\text{O}$																

3c(ii)	Answer showing:	$\text{no. of mol } C_6H_5COOH = \frac{\text{mass}}{\text{gfm}} = \frac{5.0}{122} = 0.0410\text{mol}$ $\text{no. of mol } CH_3OH = \frac{\text{mass}}{\text{gfm}} = \frac{2.5}{32} = 0.0781\text{mol}$ $C_6H_5COOH + CH_3OH \rightarrow C_6H_5COOCH_3 + H_2O$ <p style="text-align: center;"> 1mol 1mol 0.0410mol Available 0.0410mol Required </p> <p>0.0410mol CH₃OH required and 0.0781 mol CH₃OH available ∴ CH₃OH in excess and C₆H₅COOH is limiting factor as a result.</p>															
3c(iii)	12.84	500g Benzoic Acid. —————> £39.80 5g Benzoic acid. —————> £39.80 × ⁵ / ₅₀₀ = £0.3980 ∴ 3.1g methyl benzoate —————> £0.3980 100g methyl benzoate —————> £0.3980 × ¹⁰⁰ / _{3.1} = £12.84															
4a	One diagram from:	<table border="1" style="width: 100%; text-align: center;"> <tbody> <tr> <td style="width: 33%;"> $\begin{array}{ccccccc} & H & O & H & H & H & \\ & & & & & & \\ H & -C & -C & -C & -C & -C & -H \\ & & & & & & \\ & H & & H & H & H & \end{array}$ <p>pentan-2-one</p> </td> <td style="width: 33%;"> $\begin{array}{ccccccc} & H & H & O & H & H & \\ & & & & & & \\ H & -C & -C & -C & -C & -C & -H \\ & & & & & & \\ & H & H & & H & H & \end{array}$ <p>pentan-3-one</p> </td> <td style="width: 33%;"> $\begin{array}{ccccccc} & H & H & O & H & & \\ & & & & & & \\ H & -C & -C & -C & -C & -H & \\ & & & & & & \\ & & H-C-H & & H & & \\ & & & & & & \\ & & H & & & & \end{array}$ <p>3-methylbutan-2-one</p> </td> </tr> </tbody> </table>	$\begin{array}{ccccccc} & H & O & H & H & H & \\ & & & & & & \\ H & -C & -C & -C & -C & -C & -H \\ & & & & & & \\ & H & & H & H & H & \end{array}$ <p>pentan-2-one</p>	$\begin{array}{ccccccc} & H & H & O & H & H & \\ & & & & & & \\ H & -C & -C & -C & -C & -C & -H \\ & & & & & & \\ & H & H & & H & H & \end{array}$ <p>pentan-3-one</p>	$\begin{array}{ccccccc} & H & H & O & H & & \\ & & & & & & \\ H & -C & -C & -C & -C & -H & \\ & & & & & & \\ & & H-C-H & & H & & \\ & & & & & & \\ & & H & & & & \end{array}$ <p>3-methylbutan-2-one</p>												
$\begin{array}{ccccccc} & H & O & H & H & H & \\ & & & & & & \\ H & -C & -C & -C & -C & -C & -H \\ & & & & & & \\ & H & & H & H & H & \end{array}$ <p>pentan-2-one</p>	$\begin{array}{ccccccc} & H & H & O & H & H & \\ & & & & & & \\ H & -C & -C & -C & -C & -C & -H \\ & & & & & & \\ & H & H & & H & H & \end{array}$ <p>pentan-3-one</p>	$\begin{array}{ccccccc} & H & H & O & H & & \\ & & & & & & \\ H & -C & -C & -C & -C & -H & \\ & & & & & & \\ & & H-C-H & & H & & \\ & & & & & & \\ & & H & & & & \end{array}$ <p>3-methylbutan-2-one</p>															
4b	One oxidising agent from:	<table border="1" style="width: 100%; text-align: center;"> <thead> <tr> <th>Oxidising Agent</th> <th>Start Colour</th> <th>End Colour</th> </tr> </thead> <tbody> <tr> <td>Acidified Dichromate</td> <td>Orange</td> <td>Green</td> </tr> <tr> <td>Benedict's/Fehling's Solution</td> <td>Blue</td> <td>Brick Red (orange)</td> </tr> <tr> <td>Hot copper (II) oxide</td> <td>Black</td> <td>Brown</td> </tr> <tr> <td>Tollen's Reagent</td> <td>(Colourless)</td> <td>Silver mirror produced</td> </tr> </tbody> </table>	Oxidising Agent	Start Colour	End Colour	Acidified Dichromate	Orange	Green	Benedict's/Fehling's Solution	Blue	Brick Red (orange)	Hot copper (II) oxide	Black	Brown	Tollen's Reagent	(Colourless)	Silver mirror produced
Oxidising Agent	Start Colour	End Colour															
Acidified Dichromate	Orange	Green															
Benedict's/Fehling's Solution	Blue	Brick Red (orange)															
Hot copper (II) oxide	Black	Brown															
Tollen's Reagent	(Colourless)	Silver mirror produced															
4c	Permanent dipole to permanent dipole	3-methylbutanal contains a carbonyl C=O bond. There is an electronegativity difference of 1.0 within the bond making the bond polar. The δ ⁺ and δ ⁻ charges across the bond are attractive to neighbouring molecules also with a carbonyl C=O group.															
4d	It will oxidise	Aldehydes will oxidise to carboxylic acids which gives food an unpleasant taste known as rancid.															
4e(i)	Two molecules join together with the loss of water/small molecule	<table border="1" style="width: 100%;"> <tbody> <tr> <td style="width: 20%; text-align: center;">Condensation</td> <td>Two small molecules join together to form a larger molecule with the loss of a small molecules (usually water)</td> </tr> <tr> <td style="text-align: center;">Hydrolysis</td> <td>A larger molecule splits into two smaller molecules with a small molecule (usually water) added across the break point</td> </tr> </tbody> </table>	Condensation	Two small molecules join together to form a larger molecule with the loss of a small molecules (usually water)	Hydrolysis	A larger molecule splits into two smaller molecules with a small molecule (usually water) added across the break point											
Condensation	Two small molecules join together to form a larger molecule with the loss of a small molecules (usually water)																
Hydrolysis	A larger molecule splits into two smaller molecules with a small molecule (usually water) added across the break point																
4e(ii)	6-methylheptan-2-one	<table border="1" style="width: 100%;"> <tbody> <tr> <td style="width: 60%; text-align: center;"> $\begin{array}{ccccccccccc} & H & H & H & H & H & O & H & & & \\ & & & & & & & & & & \\ H & -C & -C & -C & -C & -C & -C & -C & -H & & \\ & & & & & & & & & & \\ & H & & H & H & H & & H & & & \\ & & & & & & & & & & \\ & & H-C-H & & & & & & & & \\ & & & & & & & & & & \\ & & H & & & & & & & & \end{array}$ </td> <td style="width: 40%;"> C=O Carbonyl group between two carbons ∴ -one C=O Carbonyl group on C₂ ∴ -2-one Seven carbons on main chain with functional group ∴ heptan-2-one -CH₃ methyl group on C₆ ∴ 6-methylheptan-2-one </td> </tr> </tbody> </table>	$\begin{array}{ccccccccccc} & H & H & H & H & H & O & H & & & \\ & & & & & & & & & & \\ H & -C & -C & -C & -C & -C & -C & -C & -H & & \\ & & & & & & & & & & \\ & H & & H & H & H & & H & & & \\ & & & & & & & & & & \\ & & H-C-H & & & & & & & & \\ & & & & & & & & & & \\ & & H & & & & & & & & \end{array}$	C=O Carbonyl group between two carbons ∴ -one C=O Carbonyl group on C ₂ ∴ -2-one Seven carbons on main chain with functional group ∴ heptan-2-one -CH ₃ methyl group on C ₆ ∴ 6-methylheptan-2-one													
$\begin{array}{ccccccccccc} & H & H & H & H & H & O & H & & & \\ & & & & & & & & & & \\ H & -C & -C & -C & -C & -C & -C & -C & -H & & \\ & & & & & & & & & & \\ & H & & H & H & H & & H & & & \\ & & & & & & & & & & \\ & & H-C-H & & & & & & & & \\ & & & & & & & & & & \\ & & H & & & & & & & & \end{array}$	C=O Carbonyl group between two carbons ∴ -one C=O Carbonyl group on C ₂ ∴ -2-one Seven carbons on main chain with functional group ∴ heptan-2-one -CH ₃ methyl group on C ₆ ∴ 6-methylheptan-2-one																
5	Open Question Answer to Include:	<table border="1" style="width: 100%;"> <thead> <tr> <th style="width: 33%;">3 mark answer</th> <th style="width: 33%;">2 mark answer</th> <th style="width: 33%;">1 mark answer</th> </tr> </thead> <tbody> <tr> <td>Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.</td> <td>Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.</td> <td>Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.</td> </tr> </tbody> </table>	3 mark answer	2 mark answer	1 mark answer	Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.									
3 mark answer	2 mark answer	1 mark answer															
Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.															

6a(i)	Hydrolysis	$\text{C}_{15}\text{H}_{31}-\text{C} \begin{array}{l} \text{=O} \\ \text{O}-\text{C}_{20}\text{H}_{29} \end{array} + \text{H}_2\text{O}$												
6a(ii)	$\text{C}_{20}\text{H}_{29}\text{OH}$ or $\text{C}_{20}\text{H}_{30}\text{O}$	$\text{C}_{15}\text{H}_{31}-\text{C} \begin{array}{l} \text{=O} \\ \text{OH} \end{array} + \text{H}-\text{O}-\text{C}_{20}\text{H}_{29}$												
6b(i)	Bond breaking by u.v. light	The initiation step forms free radicals by breaking covalent bonds and free radical particles are formed with unpaired electrons e.g. $\text{Cl}-\text{Cl} \rightarrow \text{Cl}^\bullet + \text{Cl}^\bullet$												
6b(ii)	Propagation	<table border="1"> <thead> <tr> <th>Step</th> <th>Reactants (before Arrow)</th> <th>Products (after Arrow)</th> </tr> </thead> <tbody> <tr> <td>Initiation</td> <td>No free radicals on Reactant Side</td> <td>Free radicals on Product Side</td> </tr> <tr> <td>Propagation</td> <td colspan="2">Free Radicals found on both sides of arrow</td> </tr> <tr> <td>Termination</td> <td>Free radicals on Reactant Side</td> <td>No free radicals on Product Side</td> </tr> </tbody> </table>	Step	Reactants (before Arrow)	Products (after Arrow)	Initiation	No free radicals on Reactant Side	Free radicals on Product Side	Propagation	Free Radicals found on both sides of arrow		Termination	Free radicals on Reactant Side	No free radicals on Product Side
Step	Reactants (before Arrow)	Products (after Arrow)												
Initiation	No free radicals on Reactant Side	Free radicals on Product Side												
Propagation	Free Radicals found on both sides of arrow													
Termination	Free radicals on Reactant Side	No free radicals on Product Side												
6b(iii)	One answer from:	<table border="1"> <tbody> <tr> <td>Can react with free radicals to form stable molecules</td> <td>Donates electrons</td> <td>Acts as a reducing agent</td> <td>Provides electrons to pair with an unpaired electron</td> </tr> </tbody> </table>	Can react with free radicals to form stable molecules	Donates electrons	Acts as a reducing agent	Provides electrons to pair with an unpaired electron								
Can react with free radicals to form stable molecules	Donates electrons	Acts as a reducing agent	Provides electrons to pair with an unpaired electron											
6c(i)	$\begin{array}{c} \text{O} \quad \text{H} \\ \parallel \quad \\ -\text{C}-\text{N}- \\ \quad \quad \end{array}$													
6c(ii)														
7a(i)	One answer from:													
7a(ii)	Sesquiterpene	Isoprene (2-methylbuta-1,3-diene) has the formula C_5H_8 . Humulene has formula $\text{C}_{15}\text{H}_{24}$ and is formed when three C_5H_8 units join together.												
7b(i)	5.345	$1 \text{ flu vaccine} \longleftrightarrow 10.69 \text{ mg squalene}$ $500000 \text{ flu vaccines} \longleftrightarrow 10.69 \text{ mg squalene} \times \frac{500000}{1}$ $= 5345000 \text{ mg squalene}$ $= 5345 \text{ g squalene}$ $= 5.345 \text{ kg squalene}$												
7b(ii)	6	<p>1 mol of H_2 will add across 1 mol $\text{C}=\text{C}$ double bonds</p> <p>6 mol of H_2 will add across 6 mol $\text{C}=\text{C}$ double bonds</p> <p>1 mol of Squalene contains 6 mol $\text{C}=\text{C}$ double bonds</p> <p>\therefore 1 mol of squalene will react with 6 mol of H_2</p>												

7c(i)	Addition or Hydration	Addition Reactions have a molecule adding across a C=C double bond or a C≡C triple bond. If the molecule adding across the C=C double bond is water then the reaction can also be described as hydration.																				
7c(ii)	Terpineol is a tertiary alcohol	Hot copper (II) oxide oxidises primary alcohols into carboxylic acids and oxidises secondary alcohols into ketones. Tertiary alcohols do not oxidise.																				
8a	286	<table border="1"> <thead> <tr> <th colspan="2">Bond Breaking Steps</th> <th colspan="2">Bond Forming Steps</th> </tr> </thead> <tbody> <tr> <td>6x C-H bonds</td> <td>6x 412kJ = 2472kJ</td> <td>1x C≡C bonds</td> <td>1x 838kJ = 838kJ</td> </tr> <tr> <td>1x C-C bond</td> <td>1x 348kJ = 348kJ</td> <td>2x C-H bonds</td> <td>2x 412kJ = 824kJ</td> </tr> <tr> <td></td> <td></td> <td>2x H-H bonds</td> <td>2x 436kJ = 872kJ</td> </tr> <tr> <td>Total bond breaking</td> <td>= 2820kJ</td> <td>Total bond Forming</td> <td>= 2534kJ</td> </tr> </tbody> </table> <p> $\Delta H = \Sigma \text{Bond enthalpies for bonds broken} - \Sigma \text{Bond enthalpies for bonds formed}$ $\Delta H = 2820 - 2534$ $\Delta H = +286 \text{ kJ mol}^{-1}$ </p>	Bond Breaking Steps		Bond Forming Steps		6x C-H bonds	6x 412kJ = 2472kJ	1x C≡C bonds	1x 838kJ = 838kJ	1x C-C bond	1x 348kJ = 348kJ	2x C-H bonds	2x 412kJ = 824kJ			2x H-H bonds	2x 436kJ = 872kJ	Total bond breaking	= 2820kJ	Total bond Forming	= 2534kJ
Bond Breaking Steps		Bond Forming Steps																				
6x C-H bonds	6x 412kJ = 2472kJ	1x C≡C bonds	1x 838kJ = 838kJ																			
1x C-C bond	1x 348kJ = 348kJ	2x C-H bonds	2x 412kJ = 824kJ																			
		2x H-H bonds	2x 436kJ = 872kJ																			
Total bond breaking	= 2820kJ	Total bond Forming	= 2534kJ																			
8b	185	$\begin{array}{l} \textcircled{1} \quad \text{C} + \text{O}_2 \rightarrow \text{CO}_2 \quad \Delta H = -394 \text{ kJ mol}^{-1} \\ \textcircled{2} \quad \text{H}_2 + \frac{1}{2}\text{O}_2 \rightarrow \text{H}_2\text{O} \quad \Delta H = -286 \text{ kJ mol}^{-1} \\ \textcircled{3} \quad \text{C}_3\text{H}_4 + 4\text{O}_2 \rightarrow 3\text{CO}_2 + 2\text{H}_2\text{O} \quad \Delta H = -1939 \text{ kJ mol}^{-1} \\ \textcircled{1} \times 3 \quad 3\text{C} + \cancel{3\text{O}_2} \rightarrow \cancel{3\text{CO}_2} \quad \Delta H = -1182 \text{ kJ} \\ \textcircled{2} \times 2 \quad 2\text{H}_2 + \cancel{\text{O}_2} \rightarrow \cancel{2\text{H}_2\text{O}} \quad \Delta H = -572 \text{ kJ} \\ \textcircled{3} \times -1 \quad \cancel{3\text{CO}_2} + \cancel{2\text{H}_2\text{O}} \rightarrow \text{C}_3\text{H}_4 + \cancel{4\text{O}_2} \quad \Delta H = +1939 \text{ kJ} \\ \text{Add} \\ \textcircled{1} + \textcircled{2} + \textcircled{3} \quad 3\text{C} + 2\text{H}_2 \rightarrow \text{C}_3\text{H}_4 \quad \Delta H = +185 \text{ kJ mol}^{-1} \end{array}$																				
8c(i)	48475	<p>gfm C₃H₄ = 40g</p> $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{1000\text{g}}{40} = 25 \text{ mol}$ <p>1mol \longleftrightarrow -1939 kJ 25mol \longleftrightarrow -1939kJ $\times \frac{25}{1}$ = 48475 kJ</p>																				
8c(ii)	13.76	<p>gfm C₃H₄ = 40g</p> $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{1\text{g}}{40\text{g mol}^{-1}} = 0.025 \text{ mol}$ $\text{C}_3\text{H}_4(\text{g}) + 4\text{O}_2(\text{g}) \longrightarrow 3\text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$ <p>1mol 4mol 0.025mol 0.1mol</p> <p>gfm O₂ = 32g</p> $\text{mass} = \text{no. of mol} \times \text{gfm} = 0.1 \times 32 = 3.2\text{g}$ <p>mass of air = 4.3 x mass of oxygen mass of air = 4.3 x 3.2 mass of air = 13.76g</p>																				
8c(iii)	Methanol and ethanol contain oxygen in their structure	Methanol and ethanol are alcohols which contain the -OH hydroxyl group. This oxygen inside the molecule means less oxygen is required from air to burn the structure fully. Ethane and propane are alkanes and these hydrocarbons have no oxygen in their structure.																				
9a(i)	Any two from:	<table border="1"> <tbody> <tr> <td>recycle (waste) gases</td> <td>use catalyst</td> <td>low/reduce energy requirements</td> </tr> <tr> <td>reactors are run at lower temperatures</td> <td>inexpensive feedstocks</td> <td>selling/using by-products</td> </tr> </tbody> </table>	recycle (waste) gases	use catalyst	low/reduce energy requirements	reactors are run at lower temperatures	inexpensive feedstocks	selling/using by-products														
recycle (waste) gases	use catalyst	low/reduce energy requirements																				
reactors are run at lower temperatures	inexpensive feedstocks	selling/using by-products																				
9a(ii)	Distillation	Distillation separates chemicals with different boiling points. Ethane-1,2-diol has a higher boiling point due to hydrogen bonding by its two hydroxyl -OH groups.																				
9b	Answer to include:	<table border="1"> <tbody> <tr> <td>1st Mark:</td> <td>Propan-1-ol has one hydroxyl -OH group Ethane-1,2-diol has two hydroxyl -OH group</td> </tr> <tr> <td>2nd Mark:</td> <td>Stronger/more hydrogen bonding in ethane-1,2-diol than propan-1-ol</td> </tr> </tbody> </table>	1 st Mark:	Propan-1-ol has one hydroxyl -OH group Ethane-1,2-diol has two hydroxyl -OH group	2 nd Mark:	Stronger/more hydrogen bonding in ethane-1,2-diol than propan-1-ol																
1 st Mark:	Propan-1-ol has one hydroxyl -OH group Ethane-1,2-diol has two hydroxyl -OH group																					
2 nd Mark:	Stronger/more hydrogen bonding in ethane-1,2-diol than propan-1-ol																					

12b(i)	126.9	$\text{C}_6\text{H}_6 + \text{H}_2\text{SO}_4 + 2\text{NaOH} \longrightarrow \text{C}_6\text{H}_5\text{OH} + \text{Na}_2\text{SO}_3 + 2\text{H}_2\text{O}$ $\begin{array}{ccc} 1\text{mol} & & 1\text{mol} \\ 78\text{g} & & 94\text{g} \\ 117\text{g} & & 94\text{g} \times \frac{117}{78} \\ & & = 141\text{g} \\ & & 141\text{kg} \end{array}$ $\% \text{Yield} = \frac{\text{Actual}}{\text{Theoretical}} \times 100 \quad \therefore \text{Actual} = \frac{\% \text{Yield} \times \text{Theoretical}}{100} = \frac{90 \times 141\text{kg}}{100} = 126.9\text{kg}$				
12b(ii)	propanone	<p>Cumene hydroperoxide \longrightarrow Phenol + X [ring]-C₃H₇O₂ \longrightarrow [ring]-OH + C₃H₆O</p> <p>C₃H₆O has two possible structures.</p> <table border="1" data-bbox="576 555 1495 786"> <thead> <tr> <th data-bbox="576 555 1038 595">Propanone</th> <th data-bbox="1038 555 1495 595">propanal</th> </tr> </thead> <tbody> <tr> <td data-bbox="576 595 1038 786"> $\begin{array}{ccccc} & \text{H} & \text{O} & \text{H} & \\ & & & & \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{H} \\ & & & & \\ & \text{H} & & \text{H} & \end{array}$ </td> <td data-bbox="1038 595 1495 786"> $\begin{array}{ccccc} & \text{H} & \text{H} & & \text{O} \\ & & & & // \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & \\ & & & & \backslash \\ & \text{H} & \text{H} & & \text{H} \end{array}$ </td> </tr> </tbody> </table> <p>The oxygen atoms in cumene hydroperoxide are attached to C₂ of three carbon chain Conclusion: X must be propanone</p>	Propanone	propanal	$\begin{array}{ccccc} & \text{H} & \text{O} & \text{H} & \\ & & & & \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{H} \\ & & & & \\ & \text{H} & & \text{H} & \end{array}$	$\begin{array}{ccccc} & \text{H} & \text{H} & & \text{O} \\ & & & & // \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & \\ & & & & \backslash \\ & \text{H} & \text{H} & & \text{H} \end{array}$
Propanone	propanal					
$\begin{array}{ccccc} & \text{H} & \text{O} & \text{H} & \\ & & & & \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{H} \\ & & & & \\ & \text{H} & & \text{H} & \end{array}$	$\begin{array}{ccccc} & \text{H} & \text{H} & & \text{O} \\ & & & & // \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & \\ & & & & \backslash \\ & \text{H} & \text{H} & & \text{H} \end{array}$					