



National
Qualifications
2024

2024 Mathematics

Higher - Paper 1

Question Paper Finalised Marking Instructions

© Scottish Qualifications Authority 2024

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.



Marking Instructions for each question

Question	Generic scheme	Illustrative scheme	Max mark
1.	<ul style="list-style-type: none"> •¹ use $m = \tan \theta$ •² evaluate exact value •³ determine equation 	<ul style="list-style-type: none"> •¹ $m = \tan 30^\circ$ •² $\frac{1}{\sqrt{3}}$ •³ eg $y = \frac{1}{\sqrt{3}}x + 4$ or $\sqrt{3}y - 4\sqrt{3} = x$ 	3

Notes:

1. Do not award •¹ for $m = \tan^{-1} 30^\circ$. However •² and •³ are still available.
2. Do not penalise the omission of a degree symbol at •¹.
3. Where candidates make no reference to a trigonometric ratio, or use an incorrect trigonometric ratio, •¹ and •² are unavailable. See Candidate A.
4. •³ is only available as a consequence of attempting to use a tan ratio. See Candidate F.
5. •³ is not available for using a gradient of 30.
6. At •³ accept any rearrangement of a candidate's equation where constant terms have been simplified.
7. Accept $y - 4 = \frac{1}{\sqrt{3}}(x)$ but not $y - 4 = \frac{1}{\sqrt{3}}(x - 0)$ for •³.

Commonly Observed Responses:

<p>Candidate A - no trig ratio</p> $m = \frac{1}{\sqrt{3}}$ • ¹ ^ • ² ✓ ₂	<p>Candidate B</p> $m = \tan \theta$ • ¹ ^	<p>Candidate C</p> $m = \tan \theta$ • ¹ ^
$y = \frac{1}{\sqrt{3}}x + 4$ • ³ ✓ ₁	$y = \frac{1}{\sqrt{3}}x + 4$ • ² ✓ • ³ ✓	$y = \sqrt{3}x + 4$ • ² x • ³ x
<p>Candidate D</p> $m = \tan \theta = 30$ • ¹ x	<p>Candidate E - no reference to m</p> $\tan 30^\circ = \frac{1}{\sqrt{3}}$ • ² ✓	<p>Candidate F - not using tan</p> $m = \sin 30^\circ$ • ¹ x
$m = \frac{1}{\sqrt{3}}$ • ² ✓ ₁	$y - 4 = \frac{1}{\sqrt{3}}(x - 0)$ • ¹ ✓	$m = \frac{1}{2}$ • ² ✓ ₂
$y = \frac{1}{\sqrt{3}}x + 4$ • ³ ✓ ₁	$y = \frac{1}{\sqrt{3}}x + 4$ • ³ ✓	$y = \frac{1}{2}x + 4$ • ³ ✓ ₂

Question		Generic scheme		Illustrative scheme	Max mark
2.	(a)		• ¹ calculate second term	• ¹ 16	1
Notes:					
1. Candidates who use $u_0 = 20$ and then calculate $u_1 = 16$ gain • ¹ .					
Commonly Observed Responses:					
	(b)	(i)	• ² communicate condition for limit to exist	• ² a limit exists as $-1 < \frac{1}{5} < 1$	1
		(ii)	• ³ know how to calculate a limit • ⁴ calculate limit	• ³ $\frac{12}{1 - \frac{1}{5}}$ or $L = \frac{1}{5}L + 12$ • ⁴ 15	2
Notes:					
2. For • ² accept: any of ' $-1 < \frac{1}{5} < 1$ ' or ' $\left \frac{1}{5}\right < 1$ ' or ' $0 < \frac{1}{5} < 1$ ' with no further comment; or statements such as: ' $\frac{1}{5}$ lies between -1 and 1 ' or ' $\frac{1}{5}$ is a proper fraction'.					
3. • ² is not available for: ' $-1 \leq \frac{1}{5} \leq 1$ ' or ' $\frac{1}{5} < 1$ ' or statements such as: 'It is between -1 and 1 .' or ' $\frac{1}{5}$ is a fraction'.					
4. Candidates who state $-1 < a < 1$ can only gain • ² if it is explicitly stated that $a = \frac{1}{5}$.					
5. Do not accept $L = \frac{b}{1-a}$ with no further working for • ³ .					
6. • ³ and • ⁴ are not available to candidates who conjecture $L = 15$ following the calculation of further terms in the sequence.					
7. For $L = 15$ with no working award 0/2.					
8. • ⁴ is only available where • ³ has been awarded.					
Commonly Observed Responses:					
Candidate A $a = \frac{1}{5}$ $-1 < a < 1$ so a limit exists			• ² ✓	Candidate B - no explicit reference to a $u_{n+1} = au_n + b$ $u_{n+1} = \frac{1}{5}u_n + 12$ $-1 < a < 1$ so a limit exists	
				• ² ^	

Question		Generic scheme	Illustrative scheme	Max mark
3.		<ul style="list-style-type: none"> •¹ start to differentiate •² complete differentiation 	<ul style="list-style-type: none"> •¹ $7(5x^2 + 3)^6 \dots$ •² $\dots \times 10x$ 	2
Notes:				
<p>1. •¹ is awarded for the appearance of $7(5x^2 + 3)^6$.</p> <p>2. For $70x(5x^2 + 3)^6$ with no working, award 2/2.</p> <p>3. Accept $7u^6$ where $u = 5x^2 + 3$ for •¹.</p> <p>4. Do not award •² where the answer includes '+c'.</p>				
Commonly Observed Responses:				
Candidate A - differentiating over two lines			Candidate B - poor notation	
$7(5x^2 + 3)^6$		• ¹ ✓	$y = (5x^2 + 3)^7$	
$7(5x^2 + 3)^6 \times 10x$		• ² ^	$y = 5x^2 + 3$	
			$\frac{dy}{dx} = 10x$	
			$\frac{dy}{dx} = 7(5x^2 + 3)^6 \times 10x$	• ¹ ✓ • ² ✓
Candidate C - poor communication			Candidate D - insufficient evidence for •¹	
$y = (5x^2 + 3)^7$			$70(5x^2 + 3)^6$	• ¹ ✗ • ² ✗
$y = 7(5x^2 + 3)^6 \times 10x$		• ¹ ✓ • ² ✓	or	
			$35(5x^2 + 3)^6$	• ¹ ✗ • ² ✗

Question		Generic scheme	Illustrative scheme	Max mark
4.		<p>Method 1</p> <ul style="list-style-type: none"> •¹ interpret ratio •² find coordinates of R 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\begin{pmatrix} 2 \\ 4 \\ -4 \end{pmatrix}, \begin{pmatrix} -2 \\ -4 \\ 4 \end{pmatrix}, \begin{pmatrix} 3 \\ 6 \\ -6 \end{pmatrix}$ or $\begin{pmatrix} -3 \\ -6 \\ 6 \end{pmatrix}$ •² $(-4, 5, -2)$ 	2
		<p>Method 2</p> <ul style="list-style-type: none"> •¹ interpret ratio •² find coordinates of R 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ eg $\overline{PR} = \frac{2}{5}\overline{PQ}, \overline{QR} = \frac{3}{5}\overline{QP}$ or $\overline{PR} = \frac{2}{3}\overline{RQ}$ •² $(-4, 5, -2)$ 	
		<p>Method 3</p> <ul style="list-style-type: none"> •¹ use section formula •² find coordinates of R 	<p>Method 3</p> <ul style="list-style-type: none"> •¹ $\frac{1}{5}(3\mathbf{p} + 2\mathbf{q})$ •² $(-4, 5, -2)$ 	

Notes:

- For $(-4, 5, -2)$ without working award 2/2.
- For $\begin{pmatrix} -4 \\ 5 \\ -2 \end{pmatrix}$ without working award 1/2.
- For $(-3, 7, -4)$ (ratio of 3:2 with working) award 1/2. See Candidate A.
- For $\begin{pmatrix} -3 \\ 7 \\ -4 \end{pmatrix}$ without working award 0/2.

Commonly Observed Responses:

Candidate A

$$\overline{PR} = \frac{3}{5}\overline{PQ}$$

•¹ x

$$R = (-3, 7, -4)$$

•² ✓₁

Candidate B

$$\frac{\overline{PR}}{\overline{RQ}} = \frac{2}{3}$$

•¹ ✓

$$3\overline{PR} = 2\overline{RQ}$$


$$3(\mathbf{r} - \mathbf{p}) = 2(\mathbf{q} - \mathbf{r})$$

$$5\mathbf{r} = 2\mathbf{q} + 3\mathbf{p}$$

Leading to correct answer of

$$R = (-4, 5, -2)$$

•² ✓

Question	Generic scheme	Illustrative scheme	Max mark
4. (continued)			
<p>Candidate C</p> $\overrightarrow{PQ} = \begin{pmatrix} 5 \\ 10 \\ -10 \end{pmatrix}$ $R = \begin{pmatrix} 2 \\ 4 \\ -4 \end{pmatrix}$ $R = \begin{pmatrix} -6 \\ 1 \\ 2 \end{pmatrix} + \begin{pmatrix} 2 \\ 4 \\ -4 \end{pmatrix}$ $R = \begin{pmatrix} -4 \\ 5 \\ -2 \end{pmatrix}$ $R(-4, 5, -2)$	<p>Candidate D</p> $\overrightarrow{PR} = \begin{pmatrix} 2 \\ 4 \\ -4 \end{pmatrix}$ $R(-8, -3, 6)$	<p>•¹ ✓</p> <p>•² ✗</p>	<p>•¹ ✓</p> <p>•² ✓</p>
<p>Candidate E - stepping out using absolute values</p>  $R(-4, 5, -2)$		<p>•¹ ✓</p> <p>•² ✓</p>	

Question		Generic scheme	Illustrative scheme	Max mark
5.		<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ equate composite function to x •² write $h(h^{-1}(x))$ in terms of $h^{-1}(x)$ •³ state inverse function 	<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $h(h^{-1}(x)) = x$ •² $2(h^{-1}(x))^3 - 7 = x$ •³ $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$ 	3
		<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ write as $y = h(x)$ and start to rearrange •² express x in terms of y •³ state inverse function 	<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $y = h(x) \Rightarrow x = h^{-1}(y)$ $y + 7 = 2x^3$ •² $x = \sqrt[3]{\frac{y+7}{2}}$ •³ $h^{-1}(y) = \sqrt[3]{\frac{y+7}{2}}$ $\Rightarrow h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$ 	
Notes:				
<p>1. In method 1, accept $2(h^{-1}(x))^3 - 7 = x$ for •¹ and •².</p> <p>2. In method 2, accept '$y + 7 = 2x^3$', without reference to $y = h(x) \Rightarrow x = h^{-1}(y)$ at •¹.</p> <p>3. In method 2, accept $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$ without reference to $h^{-1}(y)$ at •³.</p> <p>4. In method 2, beware of candidates with working where each line is not mathematically equivalent. See candidates A and B for example.</p> <p>5. At •³ stage, accept h^{-1} written in terms of any dummy variable. For example $h^{-1}(y) = \sqrt[3]{\frac{y+7}{2}}$.</p> <p>6. $y = \sqrt[3]{\frac{x+7}{2}}$ does not gain •³.</p> <p>7. $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$ with no working gains 3/3.</p>				

Question	Generic scheme	Illustrative scheme	Max mark
5. (continued)			
Commonly Observed Responses:			
<p>Candidate A</p> $h(x) = 2x^3 - 7$ $y = 2x^3 - 7$ $x = \sqrt[3]{\frac{y+7}{2}}$ $y = \sqrt[3]{\frac{x+7}{2}}$ $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$	<p>Candidate B</p> $h(x) = 2x^3 - 7$ $y = 2x^3 - 7$ $x = 2y^3 - 7$ $y = \sqrt[3]{\frac{x+7}{2}}$ $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$	<p>•¹ ✓ •² ✓</p> <p>•³ ✗</p>	<p>•¹ ✗</p> <p>•² ✓₁</p> <p>•³ ✓₁</p>
<p>Candidate C</p> $x = 2h(x)^3 - 7$ $h(x) = \sqrt[3]{\frac{x+7}{2}}$ $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$	<p>Candidate D - Method 1</p> $h(h^{-1}(x)) = 2(h^{-1}(x))^3 - 7$ $x = 2(h^{-1}(x))^3 - 7$ $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$	<p>•¹ ✗</p> <p>•² ✓₁</p> <p>•³ ✓₁</p>	<p>•² ✓</p> <p>•¹ ✓</p> <p>•³ ✓</p>
<p>Candidate E</p> $x \rightarrow x^3 \rightarrow 2x^3 \rightarrow 2x^3 - 7 = h(x)$ $\times 2 \rightarrow -7$ $\therefore +7 \rightarrow \div 2$ $\sqrt[3]{\frac{x+7}{2}}$ $h^{-1}(x) = \sqrt[3]{\frac{x+7}{2}}$	<p>Candidate F - BEWARE of incorrect notation</p> $h'(x) =$	<p>•¹ ✓</p> <p>•² ✓</p> <p>•³ ✓</p>	<p>•³ ✗</p>

Question			Generic scheme	Illustrative scheme	Max mark
6.	(a)	(i)	<ul style="list-style-type: none"> •¹ find value of $\cos p$ •² substitute into the formula for $\sin 2p$ •³ simplify answer 	<ul style="list-style-type: none"> •¹ $\cos p = \frac{2}{\sqrt{5}}$ stated or implied by •² •² $2 \times \frac{1}{\sqrt{5}} \times \frac{2}{\sqrt{5}}$ •³ $\frac{4}{5}$ 	3
		(ii)	<ul style="list-style-type: none"> •⁴ evaluate $\cos 2p$ 	<ul style="list-style-type: none"> •⁴ $\frac{3}{5}$ 	1

Notes:

1. Evidence for •¹ may appear in (a)(ii).
2. Where a candidate substitutes an incorrect value for $\cos p$ at •², •² may be awarded if the candidate has previously stated this incorrect value or it can be implied by a diagram or Pythagoras calculation. See Candidates A and B.
3. Where a candidate explicitly states a value for $\cos p$, subsequent working must follow from that value for •² to be awarded.
4. •³ is only available as a consequence of substituting into a valid formula at •².
5. Do not penalise trigonometric ratios which are less than -1 or greater than 1 throughout this question.

Commonly Observed Responses:

Candidate A - incorrect use of Pythagoras	Candidate B - no evidence of Pythagoras
$\sqrt{\sqrt{5}^2 + 1^2} = \sqrt{6}$	
$2 \times \frac{1}{\sqrt{5}} \times \frac{\sqrt{6}}{\sqrt{5}}$	$2 \times \frac{1}{\sqrt{5}} \times \frac{\sqrt{6}}{\sqrt{5}}$
$\frac{2\sqrt{6}}{5}$	$\frac{2\sqrt{6}}{5}$
Candidate C $2 \times \sin \frac{1}{\sqrt{5}} \times \cos \frac{2}{\sqrt{5}}$	
$\frac{4}{5}$	

	(b)		<ul style="list-style-type: none"> •⁵ evaluate $\sin 4p$ 	<ul style="list-style-type: none"> •⁵ $\frac{24}{25}$ 	1
--	-----	--	---	--	---

Notes:

6. •⁵ is only available for an answer expressed as a single fraction.

Commonly Observed Responses:

--	--	--	--	--	--

Question		Generic scheme	Illustrative scheme	Max mark
7.		<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ substitute for y •² write in standard quadratic form •³ determine x-coordinate •⁴ determine y-coordinate 	<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $x^2 + (2x)^2 - 14x - 8(2x) + 45 = 0$ •² $5x^2 - 30x + 45 = 0$ •³ 3 •⁴ 6 	4
		<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ substitute for x •² write in standard quadratic form •³ determine y-coordinate •⁴ determine x-coordinate 	<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $\left(\frac{y}{2}\right)^2 + y^2 - 14\left(\frac{y}{2}\right) - 8y + 45 = 0$ •² $\frac{5}{4}y^2 - 15y + 45 = 0$ •³ 6 •⁴ 3 	
		<p style="text-align: center;">Method 3</p> <ul style="list-style-type: none"> •¹ use centre and perpendicular gradient to determine equation of radius through point of contact •² substitute for y •³ determine x-coordinate •⁴ determine y-coordinate 	<p style="text-align: center;">Method 3</p> <ul style="list-style-type: none"> •¹ $x + 2y = 15$ •² $x + 2(2x) = 15$ •³ 3 •⁴ 6 	

Notes:

1. In Methods 1 and 2, treat an absence of brackets at the •¹ stage as bad form only if corrected on the next line of working.
2. In Methods 1 and 2, •¹ is only available if the '=0' appears by the •² stage.
3. In Methods 1 and 2, if a candidate arrives at an equation which is not a quadratic •³ and •⁴ are unavailable.
4. Where the quadratic obtained at •² in Methods 1 and 2, does not have repeated roots •³ and •⁴ are not available.
5. In Method 3 accept $y - 4 = -\frac{1}{2}(x - 7)$, $-\frac{1}{2} = \frac{4 - y}{7 - x}$ or equivalent for •¹.
6. In Method 3 •², •³ and •⁴ are unavailable to candidates who find the equation of any other line.
7. For (3,6) without working, award 0/4.
8. For answer of (3,6) verified in both equations, or (3,6) generated by the linear equation and verified in the equation of the circle, award 4/4.

Question	Generic scheme	Illustrative scheme	Max mark
7. (continued)			
Commonly Observed Responses:			
<p>Candidate A - substitution into the equation of the circle</p> <p>∴</p> $x = 3 \quad \bullet^3 \checkmark$ $(3)^2 + y^2 - 14(3) - 8y + 45 = 0$ $y^2 - 8y + 12 = 0$ $(y - 2)(y - 6) = 0$ $y = 6 \quad \bullet^4 \checkmark$ <p>no need to explicitly consider $y = 2$</p> <p>However,</p> <p>$(3, 6)$ and $(3, 2)$ •⁴ ✗</p>			

Question		Generic scheme	Illustrative scheme	Max mark
8.		<ul style="list-style-type: none"> •¹ use discriminant •² apply condition •³ identify roots of quadratic expression •⁴ state range with justification 	<ul style="list-style-type: none"> •¹ $(m-4)^2 - 4(1)(2m-3)$ •² $(m-4)^2 - 4(1)(2m-3) < 0$ •³ 2, 14 •⁴ $2 < m < 14$ with eg labelled sketch or table of signs 	4

Notes:

1. At •¹, treat the inconsistent use of brackets: For example $m-4^2-4(1)(2m-3)$ or $(m-4)^2-4 \times 1 \times 2m-3$ as bad form only if the candidate deals with the unbracketed terms correctly in the next line of working.
2. Where candidates express a , b and c in terms of m , and then state $b^2-4ac < 0$, award •².
3. If candidates have the condition 'discriminant > 0 ', 'discriminant ≤ 0 ' or 'discriminant ≥ 0 ', then •² is lost but •³ and •⁴ are available.
4. Ignore the appearance of $b^2-4ac = 0$ where the correct condition has subsequently been applied.
5. If candidates only work with the condition 'discriminant = 0', then •² and •⁴ are unavailable.
6. Accept the appearance of 2 and 14 within inequalities for •³.
7. At •⁴ accept " $m > 2$ and $m < 14$ " or " $m > 2, m < 14$ " together with the required justification.
8. For the appearance of x in any expression of the discriminant, no further marks are available.

Commonly Observed Responses:


Candidate A - no expressions for a , b and c

No real roots $b^2-4ac < 0$

$$m^2 - 16m + 28 = 0$$

$$m = 2, m = 14$$

$$2 < m < 14$$



- ¹ ✓
- ³ ✓
- ² ✓ •⁴ ✓

In this case •² is only available where •⁴ is awarded

Candidate B


$$(m-4)^2 - 4(1)(2m-3)$$

$$m^2 - 16m + 28 = 0$$

$$m = 2, m = 14$$

$$b^2 - 4ac < 0$$

$$2 < m < 14$$



- ¹ ✓
- ³ ✓
- ² ✓ •⁴ ✓

In this case •² is only available where •⁴ is awarded

Question	Generic scheme	Illustrative scheme	Max mark
8. (continued)			
Candidate C $(m-4)^2 - 4(1)(2m-3)$ $b^2 - 4ac = 0$ $m^2 - 16m + 28 = 0$ $m = 2, m = 14$ $m^2 - 16m + 28 < 0$ $2 < m < 14$	$\bullet^1 \checkmark$ $\bullet^3 \checkmark$ $\bullet^2 \checkmark$ $\bullet^4 \checkmark$	Candidate D $(m-4)^2 - 4(1)(2m-3)$ $m^2 - 16m + 28 = 0$ $m = 2, m = 14$ $2 < m < 14$	$\bullet^1 \checkmark$ $\bullet^2 \times$ $\bullet^3 \checkmark$ $\bullet^4 \checkmark_2$
Candidate E - not solving a quadratic $m - 4^2 - 4(1)(2m - 3) < 0$ $-7m - 4 < 0$ $m > -\frac{4}{7}$	$\bullet^1 \times \bullet^2 \checkmark \bullet^3 \times$ $\bullet^4 \checkmark_2$		

Question		Generic scheme	Illustrative scheme	Max mark
9.		<p>Method 1</p> <ul style="list-style-type: none"> •¹ apply $\log_a x + \log_a y = \log_a xy$ •² apply $m \log_a x = \log_a x^m$ •³ apply $\log_a x - \log_a y = \log_a \frac{x}{y}$ and express in required form 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\log_a (5 \times 80) \dots$ stated or implied by •³ •² $\dots - \log_a 10^2$ stated or implied by •³ •³ $\log_a 4$ 	3
		<p>Method 2</p> <ul style="list-style-type: none"> •¹ apply $m \log_a x = \log_a x^m$ •² apply $\log_a x - \log_a y = \log_a \frac{x}{y}$ •³ apply $\log_a x + \log_a y = \log_a xy$ and express in required form 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $\dots - \log_a 10^2$ stated or implied by •³ •² $\dots + \log_a \left(\frac{80}{10^2} \right)$ stated or implied by •³ •³ $\log_a 4$ 	

Notes:

1. Where an error at the •¹ or •² stage leads to a non-integer value for k , •³ is still available.
2. Each line of working must be equivalent to the line above within a valid strategy. See commonly observed responses.
3. Where candidates apply the laws of logarithms in the incorrect order see Candidates A and B.
4. Where candidates do not consider the '2', a maximum of 1/3 is available. See Candidate C.
5. Do not penalise the omission of the base of the logarithm.
6. Correct answer with no working, award 3/3.
7. Where candidates form an invalid equation, •¹ and •² may only be awarded for working with $\log_a 5 + \log_a 80 - 2 \log_a 10$ on one side of the equation; •³ is not available.

Commonly Observed Responses:

Candidate A

$$\log_a 5 + 2 \log_a \left(\frac{80}{10} \right)$$

$$2 \log_a \left(\frac{5 \times 80}{10} \right)$$

$$\log_a (40)^2$$

$$\log_a 1600$$

Award 1/3

Candidate B

$$\log_a 400 - 2 \log_a 10$$

$$2 \log_a \left(\frac{400}{10} \right)$$

$$\log_a (40)^2$$

$$\log_a 1600$$

Award 2/3

Candidate C - ignoring the 2

$$\log_a 5 + \log_a 80 - 2 \log_a 10$$

$$\log_a 5 + \log_a \frac{80}{10}$$

$$\log_a 40$$

Award 1/3

Question		Generic scheme	Illustrative scheme	Max mark
10.	(a)	<p>•¹ use 1 in synthetic division or in evaluation of quartic</p> <p>•² complete division/evaluation and interpret result</p>	<p>•¹</p> $\begin{array}{r rrrrr} 1 & 2 & 3 & -4 & -3 & 2 \\ & & & & & \square \end{array}$ <p>or $2 \times (1)^4 + 3 \times (1)^3 - 4 \times (1)^2 - 3 \times (1) + 2$</p> <p>•²</p> $\begin{array}{r rrrrr} 1 & 2 & 3 & -4 & -3 & 2 \\ & & 2 & 5 & 1 & -2 \\ \hline & 2 & 5 & 1 & -2 & \boxed{0} \end{array}$ <p>Remainder = 0 $\therefore (x-1)$ is a factor</p> <p>or</p> <p>$f(1) = 0 \therefore (x-1)$ is a factor</p>	2

Notes:

- Communication at •² must be consistent with working at that stage i.e. a candidate's working must arrive legitimately at 0 before •² can be awarded.
- Accept any of the following for •²:
 - ' $f(1) = 0$ so $(x-1)$ is a factor'
 - 'since remainder = 0, it is a factor'
 - the '0' from any method linked to the word 'factor' by 'so', 'hence', \therefore , \rightarrow , \Rightarrow etc.
- Do not accept any of the following for •²:
 - double underlining the '0' or boxing the '0' without comment
 - ' $x=1$ is a factor', ' \dots is a root'
 - the word 'factor' only, with no link.

Commonly Observed Responses:

Candidate A - grid method

$$\begin{array}{r} 2x^3 \\ x \quad \begin{array}{|c|c|c|c|} \hline 2x^4 & 5x^3 & & \\ \hline -2x^3 & & & \\ \hline \end{array} \end{array} \quad \bullet^1 \checkmark$$

$$\begin{array}{r} 2x^3 \quad 5x^2 \quad x \quad -2 \\ x \quad \begin{array}{|c|c|c|c|} \hline 2x^4 & 5x^3 & x^2 & -2x \\ \hline -2x^3 & -5x^2 & -x & 2 \\ \hline \end{array} \end{array}$$

'with no remainder'

$\therefore (x-1)$ is a factor •² ✓

Candidate B - grid method

$$\begin{array}{r} 2x^3 \\ x \quad \begin{array}{|c|c|c|c|} \hline 2x^4 & 5x^3 & & \\ \hline -2x^3 & & & \\ \hline \end{array} \end{array} \quad \bullet^1 \checkmark$$

$$\begin{array}{r} 2x^3 \quad 5x^2 \quad x \quad -2 \\ x \quad \begin{array}{|c|c|c|c|} \hline 2x^4 & 5x^3 & x^2 & -2x \\ \hline -2x^3 & -5x^2 & -x & 2 \\ \hline \end{array} \end{array}$$

$\therefore (x-1)(2x^3 + 5x^2 + x - 2) = 2x^4 + 3x^3 - 4x^2 - 3x + 2$

$\therefore (x-1)$ is a factor •² ✓

Question		Generic scheme	Illustrative scheme	Max mark
10.	(b)	<ul style="list-style-type: none"> •³ identify cubic and attempt to factorise •⁴ find second factor •⁵ identify quadratic •⁶ complete factorisation 	<ul style="list-style-type: none"> •³ eg $\begin{array}{r rrrr} -1 & 2 & 5 & 1 & -2 \\ & & -2 & -3 & \\ \hline & 2 & 3 & & \end{array}$ or $\begin{array}{r rrrr} -2 & 2 & 5 & 1 & -2 \\ & & -4 & -2 & \\ \hline & 2 & 1 & & \end{array}$ •⁴ eg $\begin{array}{r rrrr} -1 & 2 & 5 & 1 & -2 \\ & & -2 & -3 & 2 \\ \hline & 2 & 3 & -2 & 0 \end{array}$ leading to $(x+1)$ or $\begin{array}{r rrrr} -2 & 2 & 5 & 1 & -2 \\ & & -4 & -2 & 2 \\ \hline & 2 & 1 & -1 & 0 \end{array}$ leading to $(x+2)$ •⁵ $2x^2 + 3x - 2$ or $2x^2 + x - 1$ •⁶ $(x-1)(x+1)(2x-1)(x+2)$ stated explicitly 	4

Notes:

4. Ignore the appearance of ' $= 0$ '.
5. Candidates who arrive at $(x-1)(x+1)(2x^2 + 3x - 2)$ or $(x-1)(x+2)(2x^2 + x - 1)$ by using algebraic long division or by inspection, gain •³, •⁴ and •⁵.
6. Where a candidate only identifies additional factors from a quartic, only •⁴ is available.
7. •³ and •⁴ may be awarded for applications of synthetic division even when previous trials contain errors. •⁵ and •⁶ are available.

Question	Generic scheme	Illustrative scheme	Max mark																																																																												
10. (b) (continued)																																																																															
Commonly Observed Responses:																																																																															
<p>Candidate C - grid method</p> <p>(a)</p> <table border="1" style="margin-left: 40px;"> <tr><td></td><td>$2x^3$</td><td>$5x^2$</td><td>x</td><td>-2</td></tr> <tr><td>x</td><td>$2x^4$</td><td>$5x^3$</td><td>x^2</td><td>$-2x$</td></tr> <tr><td>-1</td><td>$-2x^3$</td><td>$-5x^2$</td><td>$-x$</td><td>2</td></tr> </table> <p>(b)</p> <table border="1" style="margin-left: 40px;"> <tr><td></td><td>$2x^2$</td><td>...</td><td>...</td></tr> <tr><td>x</td><td>$2x^3$</td><td>...</td><td>...</td></tr> <tr><td>...</td><td>...</td><td>...</td><td>...</td></tr> </table> <p>•³ ✓</p> <p>•³ is awarded for evidence of the cubic expression (which may be in the grid from part (a)) AND the terms in the diagonal boxes summing to the second and third terms in the cubic respectively.</p> <table border="1" style="margin-left: 40px;"> <tr><td></td><td>$2x^2$</td><td>$3x$</td><td>-2</td></tr> <tr><td>x</td><td>$2x^3$</td><td>$3x^2$</td><td>$-2x$</td></tr> <tr><td>$+1$</td><td>$2x^2$</td><td>$3x$</td><td>-2</td></tr> </table> <p>•⁴ ✓</p> <p>$2x^2 + 3x - 2$ •⁵ ✓</p> <p>$(x-1)(x+1)(2x-1)(x+2)$ •⁶ ✓</p>		$2x^3$	$5x^2$	x	-2	x	$2x^4$	$5x^3$	x^2	$-2x$	-1	$-2x^3$	$-5x^2$	$-x$	2		$2x^2$	x	$2x^3$		$2x^2$	$3x$	-2	x	$2x^3$	$3x^2$	$-2x$	$+1$	$2x^2$	$3x$	-2	<p>Candidate D - grid method</p> <p>(a)</p> <table border="1" style="margin-left: 40px;"> <tr><td></td><td>$2x^3$</td><td>$5x^2$</td><td>x</td><td>-2</td></tr> <tr><td>x</td><td>$2x^4$</td><td>$5x^3$</td><td>x^2</td><td>$-2x$</td></tr> <tr><td>-1</td><td>$-2x^3$</td><td>$-5x^2$</td><td>$-x$</td><td>2</td></tr> </table> <p>(b)</p> <table border="1" style="margin-left: 40px;"> <tr><td></td><td>$2x^2$</td><td>...</td><td>...</td></tr> <tr><td>x</td><td>$2x^3$</td><td>...</td><td>...</td></tr> <tr><td>...</td><td>...</td><td>...</td><td>...</td></tr> </table> <p>•³ ✓</p> <p>•³ is awarded for evidence of the cubic expression (which may be in the grid from part (a)) AND the terms in the diagonal boxes summing to the second and third terms in the cubic respectively.</p> <table border="1" style="margin-left: 40px;"> <tr><td></td><td>$2x^2$</td><td>x</td><td>-1</td></tr> <tr><td>x</td><td>$2x^3$</td><td>x^2</td><td>$-x$</td></tr> <tr><td>$+2$</td><td>$4x^2$</td><td>$2x$</td><td>-2</td></tr> </table> <p>•⁴ ✓</p> <p>$2x^2 + x - 1$ •⁵ ✓</p> <p>$(x-1)(x+2)(x+1)(2x-1)$ •⁶ ✓</p>		$2x^3$	$5x^2$	x	-2	x	$2x^4$	$5x^3$	x^2	$-2x$	-1	$-2x^3$	$-5x^2$	$-x$	2		$2x^2$	x	$2x^3$		$2x^2$	x	-1	x	$2x^3$	x^2	$-x$	$+2$	$4x^2$	$2x$	-2
	$2x^3$	$5x^2$	x	-2																																																																											
x	$2x^4$	$5x^3$	x^2	$-2x$																																																																											
-1	$-2x^3$	$-5x^2$	$-x$	2																																																																											
	$2x^2$																																																																												
x	$2x^3$																																																																												
...																																																																												
	$2x^2$	$3x$	-2																																																																												
x	$2x^3$	$3x^2$	$-2x$																																																																												
$+1$	$2x^2$	$3x$	-2																																																																												
	$2x^3$	$5x^2$	x	-2																																																																											
x	$2x^4$	$5x^3$	x^2	$-2x$																																																																											
-1	$-2x^3$	$-5x^2$	$-x$	2																																																																											
	$2x^2$																																																																												
x	$2x^3$																																																																												
...																																																																												
	$2x^2$	x	-1																																																																												
x	$2x^3$	x^2	$-x$																																																																												
$+2$	$4x^2$	$2x$	-2																																																																												
<p>Candidate E</p> <table border="1" style="margin-left: 40px;"> <tr><td>$\frac{1}{2}$</td><td>2</td><td>5</td><td>1</td><td>-2</td></tr> <tr><td></td><td></td><td>1</td><td>3</td><td>2</td></tr> <tr><td></td><td>2</td><td>6</td><td>4</td><td>0</td></tr> </table> <p>•³ ✓ •⁴ ✓</p> <p>•⁵ ✓</p> <p>$(x-\frac{1}{2})(2x^2+6x+4)$</p> <p>$(2x-1)(x^2+3x+2)$</p> <p>$(x-1)(2x-1)(x+1)(x+2)$ •⁶ ✓</p>	$\frac{1}{2}$	2	5	1	-2			1	3	2		2	6	4	0	<p>Candidate F</p> <table border="1" style="margin-left: 40px;"> <tr><td>$\frac{1}{2}$</td><td>2</td><td>5</td><td>1</td><td>-2</td></tr> <tr><td></td><td></td><td>1</td><td>3</td><td>2</td></tr> <tr><td></td><td>2</td><td>6</td><td>4</td><td>0</td></tr> </table> <p>•³ ✓ •⁴ ✓</p> <p>•⁵ ✓</p> <p>$(x-\frac{1}{2})(2x^2+6x+4)$</p> <p>$(x-\frac{1}{2})(2x+2)(x+2)$</p> <p>$(x-1)(x-\frac{1}{2})(2x+2)(x+2)$ •⁶ ^</p>	$\frac{1}{2}$	2	5	1	-2			1	3	2		2	6	4	0																																																
$\frac{1}{2}$	2	5	1	-2																																																																											
		1	3	2																																																																											
	2	6	4	0																																																																											
$\frac{1}{2}$	2	5	1	-2																																																																											
		1	3	2																																																																											
	2	6	4	0																																																																											

Question		Generic scheme	Illustrative scheme	Max mark
11.	(a)	<ul style="list-style-type: none"> •¹ use compound angle formula •² compare coefficients •³ process for k •⁴ process for a and express in required form 	<ul style="list-style-type: none"> •¹ $k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$ stated explicitly •² $k \cos a^\circ = 1, k \sin a^\circ = \sqrt{3}$ stated explicitly •³ $k = 2$ •⁴ $2 \cos(x - 60)^\circ$ 	4


Notes:

1. Accept $k(\cos x^\circ \cos a^\circ + \sin x^\circ \sin a^\circ)$ for •¹. Treat $k \cos x^\circ \cos a^\circ + \sin x^\circ \sin a^\circ$ as bad form only if the equations at the •² stage both contain k .
2. Do not penalise the omission of degree signs.
3. $2 \cos x^\circ \cos a^\circ + 2 \sin x^\circ \sin a^\circ$ or $2(\cos x^\circ \cos a^\circ + \sin x^\circ \sin a^\circ)$ is acceptable for •¹ and •³.
4. •² is not available for $k \cos x^\circ = 1, k \sin x^\circ = \sqrt{3}$, however •⁴ may still be gained- see Candidate E
5. •³ is only available for a single value of $k, k > 0$.
6. •³ is not available to candidates who work with $\sqrt{4}$ throughout parts (a) and (b) without explicitly simplifying at any stage. •⁴ is still available.
7. •⁴ is not available for a value of a given in radians.
8. Candidates may use any form of the wave function for •¹, •² and •³. However, •⁴ is only available if the wave is interpreted in the form $k \cos(x - a)^\circ$.
9. Evidence for •⁴ may not appear until part (b).

Commonly Observed Responses:

Candidate A		Candidate B - inconsistency		Candidate C	
	• ¹ ^	$k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$	• ¹ ✓	$\cos x^\circ \cos a^\circ + \sin x^\circ \sin a^\circ$	• ¹ ✗
$2 \cos a^\circ = 1$		$\cos a^\circ = 1$		$\cos a^\circ = 1$	
$2 \sin a^\circ = \sqrt{3}$	• ² ✓ • ³ ✓	$\sin a^\circ = \sqrt{3}$	• ² ✗	$\sin a^\circ = \sqrt{3}$	• ² ✓ ₂
		$k = 2$		$k = 2$	• ³ ✓
$\tan a^\circ = \sqrt{3}$		$\tan a^\circ = \sqrt{3}$		$\tan a^\circ = \sqrt{3}$	
$a = 60$		$a = 60$		$a = 60$	
$2 \cos(x - 60)^\circ$	• ⁴ ✓	$2 \cos(x - 60)^\circ$	• ³ ✓ • ⁴ ✗	$2 \cos(x - 60)^\circ$	• ⁴ ✗

Question	Generic scheme	Illustrative scheme	Max mark
11. (a) (continued)			
<p>Candidate D - errors at ●² $k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$ ●¹✓ $k \cos a^\circ = \sqrt{3}$ $k \sin a^\circ = 1$ ●² ✗ $\tan a^\circ = \frac{1}{\sqrt{3}}$ $a = 30$ $2 \cos(x - 30)^\circ$ ●³✓ ●⁴✓₁</p>	<p>Candidate E - use of x at ●² $k \cos x^\circ \cos a^\circ + k \sin x^\circ \sin a^\circ$ ●¹✓ $k \cos x^\circ = 1$ $k \sin x^\circ = \sqrt{3}$ ●² ✗ $\tan x^\circ = \sqrt{3}$ $x = 60$ $2 \cos(x - 60)^\circ$ ●³✓ ●⁴✓₁</p>	<p>Candidate F $k \sin A \cos B + k \cos A \sin B$ ●¹ ✗ $k \cos A = 1$ $k \sin A = \sqrt{3}$ ●² ✗ $\tan A = \sqrt{3}$ $2 \cos(x - 60)^\circ$ ●³✓ ●⁴✓₁</p>	

Question		Generic scheme	Illustrative scheme	Max mark
11.	(b)	<ul style="list-style-type: none"> •⁵ exactly two roots identifiable from graph •⁶ coordinates of exactly two turning points identifiable from graph •⁷ y-intercept and value of y at $x = 360$ identifiable from graph 	<ul style="list-style-type: none"> •⁵ (150,0) and (330,0) •⁶ (60,2) and (240,-2) •⁷ 1 	3

Notes:

10. •⁵, •⁶ and •⁷ are only available for attempting to draw a “cosine” graph with a period of 360° .
11. Ignore any part of a graph drawn outwith $0 \leq x \leq 360$.
12. Vertical marking is not applicable to •⁵ and •⁶.
13. Candidate’s sketch in (b) must be consistent with the equation obtained in (a), see also Candidates G and H.
14. For any incorrect horizontal translation of the graph of the wave function arrived at in part (a) only •⁶ is available.


Commonly Observed Responses:

Candidate G - incorrect translation

- (a) $2 \cos(x - 60)^\circ$ - correct equation
- (b) Incorrect translation:
Sketch of $2 \cos(x + 60)^\circ$
only •⁶ is available

Candidate H - incorrect equation

- (a) $2 \cos(x + 60)^\circ$ - incorrect equation
- (b) Sketch of $2 \cos(x + 60)^\circ$
all 3 marks available



Question	Generic scheme	Illustrative scheme	Max mark
12.	<ul style="list-style-type: none"> •¹ write in differentiable form •² differentiate •³ solve for $a^{-\frac{2}{3}}$ or $a^{\frac{2}{3}}$ •⁴ solve for a 	<ul style="list-style-type: none"> •¹ $12x^{\frac{1}{3}}$ stated or implied by •² •² $12 \times \frac{1}{3} \times x^{-\frac{2}{3}}$ •³ $a^{-\frac{2}{3}} = \frac{1}{4}$ or $a^{\frac{2}{3}} = 4$ •⁴ $a = 8$ 	4

Notes:

1. •² is only available for differentiating a term with a fractional index.
2. Where candidates attempt to integrate or make no attempt to differentiate, only •¹ is available.
3. Accept $x^{-\frac{2}{3}} = \frac{1}{4}$ or $x^{\frac{2}{3}} = 4$ at •³. See Candidates A and B.
4. •⁴ is only available where the expression at •² is of the form $kx^{-\frac{m}{n}}$ where $m \neq 1$.
5. Do not penalise the inclusion of -8 at •⁴.

Commonly Observed Responses:

<p>Candidate A - working in terms of x throughout</p> <p>... •¹ ✓ •² ✓</p> <p>$x^{-\frac{2}{3}} = \frac{1}{4}$ •³ ✓</p> <p>$x = 8$ •⁴ ✗</p>	<p>Candidate B</p> <p>... •¹ ✓ •² ✓</p> <p>$x^{-\frac{2}{3}} = \frac{1}{4}$ •³ ✓</p> <p>$(x = 8)$</p> <p>$a = 8$ •⁴ ✓</p>
<p>Candidate C</p> <p>$f(x) = 12x^{\frac{3}{2}}$ •¹ ✗</p> <p>$f'(x) = 18x^{\frac{1}{2}}$ •² ✓₁</p> <p>$a^{\frac{1}{2}} = \frac{1}{18}$ •³ ✓₁</p> <p>$a = \frac{1}{324}$ •⁴ ✓₂</p>	<p>Candidate D - partly differentiated</p> <p>$f(x) = 12x^{\frac{1}{3}}$ •¹ ✓</p> <p>$f'(x) = 12 \times \frac{1}{3} x^{-\frac{2}{3}}$ •² ✗</p> <p>$1 = 4a^{\frac{4}{3}}$</p> <p>$\frac{1}{4} = a^{\frac{4}{3}}$ •³ ✓₁</p> <p>$a = \frac{1}{\sqrt[3]{8}}$ •⁴ ✓₂</p>

Question		Generic scheme	Illustrative scheme	Max mark
13.	(a)	<ul style="list-style-type: none"> •¹ find midpoint of PQ •² find gradient of PQ •³ find perpendicular gradient •⁴ find equation of perpendicular bisector 	<ul style="list-style-type: none"> •¹ (5,6) •² -4 or $-\frac{8}{2}$ •³ $\frac{1}{4}$ •⁴ $4y = x + 19$ 	4
Notes:				
<ol style="list-style-type: none"> 1. •⁴ is only available as a consequence of using a perpendicular gradient and a mid-point. 2. The gradient of the perpendicular bisector must appear in fully simplified form at •³ or •⁴ stage for •³ to be awarded. 3. At •⁴ accept $4y - x = 19$, $4y - x - 19 = 0$, or any other rearrangement of the equation where the constant terms have been simplified. 				
Commonly Observed Responses:				
	(b)	<ul style="list-style-type: none"> •⁵ identify x-coordinate of centre •⁶ find y-coordinate of centre •⁷ find radius •⁸ state equation of circle 	<ul style="list-style-type: none"> •⁵ 9 •⁶ 7 •⁷ $\sqrt{34}$ •⁸ $(x-9)^2 + (y-7)^2 = 34$ 	4
Notes:				
<ol style="list-style-type: none"> 4. Do not accept “centre = (9,2)” as evidence of •⁵. 5. Where candidates use PQ, QR or PR as the diameter of the circle no marks are available. 6. •⁷ and •⁸ are only available as a consequence of using the point of intersection of two perpendicular bisectors and a point on the circumference of the circle. 7. Accept $r^2 = 34$ for •⁷. 8. $(x-9)^2 + (y-7)^2 = (\sqrt{34})^2$ does not gain •⁸. 				
Commonly Observed Responses:				
Candidate A - horizontal line through midpoint of PQ		Candidate B - perpendicular bisector of PR		
Centre = (9,6)		Perpendicular bisector of PR: $y = x - 2$		
Radius = 5		Centre = (9,7)		
Equation: $(x-9)^2 + (y-6)^2 = 25$		∴		
• ⁵ ✓ • ⁶ ✗		• ⁵ ✓ • ⁶ ✓		
• ⁷ ✗				
• ⁸ ✗				

[END OF MARKING INSTRUCTIONS]



National
Qualifications
2024

2024 Mathematics

Higher - Paper 2

Question Paper Finalised Marking Instructions

© Scottish Qualifications Authority 2024

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments.

The information in this document may be reproduced in support of SQA qualifications only on a non-commercial basis. If it is reproduced, SQA must be clearly acknowledged as the source. If it is to be reproduced for any other purpose, written permission must be obtained from permissions@sqa.org.uk.



Question		Generic scheme	Illustrative scheme	Max mark
1.	(a)	<ul style="list-style-type: none"> •¹ determine midpoint of AC •² determine gradient of median •³ find equation of median 	<ul style="list-style-type: none"> •¹ (4,4) •² 2 or $\frac{10}{5}$ •³ $y = 2x - 4$ 	3

Notes:

1. •² is only available to candidates who use a midpoint to find a gradient.
2. •³ is only available as a consequence of using a 'midpoint' of AC and the point B
3. At •³ accept any arrangement of a candidate's equation where the constant terms have been simplified.
4. •³ is not available as a consequence of using a perpendicular gradient.

Commonly Observed Responses:

<p>Candidate A - perpendicular bisector of AC Midpoint = (4,4) •¹ ✓</p> <p>$m_{AC} = -\frac{4}{7} \Rightarrow m_{\perp} = \frac{7}{4}$ •² ✗</p> <p>$4y = 7x - 12$ •³ ✓₂</p> <p>For other perpendicular bisectors award 0/3</p>	<p>Candidate B - altitude through B</p> <p>$m_{AC} = -\frac{4}{7}$ •¹ ^</p> <p>$m_{\perp} = \frac{7}{4}$ •² ✗</p> <p>$4y = 7x - 17$ •³ ✓₂</p>
<p>Candidate C - median through A midpoint BC = (5, -3) •¹ ✗</p> <p>$m_{AM} = -\frac{11}{8}$ •² ✓₁</p> <p>$8y = -11x + 31$ •³ ✓₂</p>	<p>Candidate D - median through C midpoint AB (-2, 1) •¹ ✗</p> <p>$m_{CM} = -\frac{1}{13}$ •² ✓₁</p> <p>$13y = -x + 11$ •³ ✓₂</p>

	(b)	<ul style="list-style-type: none"> •⁴ determine gradient of BC •⁵ determine gradient of L •⁶ find equation of L 	<ul style="list-style-type: none"> •⁴ $\frac{6}{12}$ •⁵ $-\frac{12}{6}$ •⁶ $y = -2x + 22$ 	3
--	-----	---	--	---

Notes:

5. •⁶ is only available as a consequence of using a perpendicular gradient and C.
6. At •⁶ accept any arrangement of a candidate's equation where the constant terms have been simplified.

Commonly Observed Responses:

<p>Candidate E - altitude through C</p> <p>$m_{AB} = -7$ •⁴ ✗</p> <p>$m_{\perp} = \frac{1}{7}$ •⁵ ✓₁</p> <p>$y = \frac{1}{7}(x - 11)$ •⁶ ✓₁</p>	
---	--

Question		Generic scheme	Illustrative scheme	Max mark
1.	(c)	<ul style="list-style-type: none"> •⁷ determine x-coordinate •⁸ determine y-coordinate 	<ul style="list-style-type: none"> •⁷ 6.5 or $\frac{13}{2}$ •⁸ 9 	2
Notes:				
7. For $\left(\frac{26}{4}, 9\right)$ award 1/2.				
Commonly Observed Responses:				
Candidate F - rounding decimals (a) $4y = 5x - 19$ (b) $y = -2x + 22$ (c) $x = \frac{107}{13} = 8.2$ $y = 5.6$			<ul style="list-style-type: none"> •⁷ ✓₁ •⁸ ✓₁ 	

Question		Generic scheme	Illustrative scheme	Max mark
2.		<ul style="list-style-type: none"> •¹ find y-coordinate •² write in differentiable form •³ differentiate •⁴ find gradient of tangent •⁵ determine equation of tangent 	<ul style="list-style-type: none"> •¹ 1 •² $8x^{-3}$ •³ $8 \times (-3)x^{-4}$ •⁴ $-\frac{3}{2}$ •⁵ $3x + 2y = 8$ 	5

Notes:

1. Only •¹ and •² are available to candidates who integrate. However, see Candidates E and F.
2. $8 \times (-3)x^{-4}$ without previous working gains •² and •³.
3. •³ is only available for differentiating a negative power. •⁴ and •⁵ are still available.
4. •⁴ is not available for $y = -\frac{3}{2}$. However, where $-\frac{3}{2}$ is then used as the gradient of the straight line, •⁴ may be awarded - see Candidates A, B and C.
5. •⁵ is only available where candidates attempt to find the gradient by substituting into their derivative.
6. •⁵ is not available as a consequence of using a perpendicular gradient.
7. Where $x = 2$ has not been used to determine the y-coordinate, •⁵ is not available.

Commonly Observed Responses:

<p>Candidate A - incorrect notation</p> $y = 1$ $y = 8x^{-3}$ $y = -24x^{-4}$ $y = -\frac{3}{2}$ $3x + 2y = 8$ <ul style="list-style-type: none"> •¹ ✓ - BoD •² ✓ •³ ✓ •⁴ ✓ - BoD •⁵ ✓ 	<p>Candidate B - use of values in equation</p> $y = 1$ $y = 8x^{-3}$ $\frac{dy}{dx} = 8 \times (-3)x^{-4}$ $\frac{dy}{dx} = -\frac{3}{2}$ $y = -\frac{3}{2}$ $3x + 2y = 8$ <ul style="list-style-type: none"> •¹ ✓ - BoD •² ✓ •³ ✓ •⁴ ✓ •⁵ ✓
<p>Candidate C - incorrect notation</p> $y = 1$ $y = 8x^{-3}$ $\frac{dy}{dx} = 8 \times (-3)x^{-4}$ $y = -\frac{3}{2}$ <ul style="list-style-type: none"> •¹ ✓ - BoD •² ✓ •³ ✓ •⁴ ✗ <p>Evidence for •⁴ would need to appear in the equation of the line</p>	<p>Candidate D</p> $y = 1$ $y = 8x^{-3}$ $\frac{dy}{dx} = 8 \times (-3)x^{-4} = 0$ $8 \times (-3)(2)^{-4} = 0$ $m = -\frac{3}{2}$ $3x + 2y = 8$ <ul style="list-style-type: none"> •¹ ✓ •² ✓ •³ ✓ •⁴ ✗ •⁵ ✓₁

Question	Generic scheme	Illustrative scheme	Max mark
2. (continued)			
Candidate E - integrating in part $y = 1$ $y = 8x^{-3}$ $\frac{dy}{dx} = -24x^{-2}$ $\frac{dy}{dx} = -6$ $y = -6x + 13$	Candidate F - appearance of $+c$ $y = 1$ $y = 8x^{-3}$ $\frac{dy}{dx} = -24x^{-4} + c$	 	


Question		Generic scheme	Illustrative scheme	Max mark	
3.	(a)	<ul style="list-style-type: none"> •¹ find \vec{ED} •² find \vec{EF} 	<ul style="list-style-type: none"> •¹ $\begin{pmatrix} 1 \\ -4 \\ 6 \end{pmatrix}$ •² $\begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$ 	2	
Notes:					
1. For candidates who find both \vec{DE} and \vec{FE} correctly, award 1/2.					
2. Accept vectors written horizontally.					
Commonly Observed Responses:					
	(b)	(i)	• ³ evaluate $\vec{ED} \cdot \vec{EF}$	• ³ 16	1
		(ii)	<ul style="list-style-type: none"> •⁴ evaluate \vec{ED} •⁵ evaluate \vec{EF} •⁶ substitute into formula for scalar product •⁷ calculate angle 	<ul style="list-style-type: none"> •⁴ $\sqrt{53}$ •⁵ $\sqrt{14}$ •⁶ $\cos DEF = \frac{16}{\sqrt{53} \times \sqrt{14}}$ or $\sqrt{53} \times \sqrt{14} \times \cos DEF = 16$ •⁷ $54.028\dots^\circ$ or $0.942\dots$ radians 	4

Question	Generic scheme	Illustrative scheme	Max mark
3. (b) (continued)			
Notes:			
<p>3. Do not penalise candidates who treat negative signs with a lack of rigour when calculating a magnitude. For example accept $\sqrt{1^2 + 4^2 + 6^2} = \sqrt{53}$ or $\sqrt{1^2 + -4^2 + 6^2} = \sqrt{53}$ for \bullet^4. However, do not accept $\sqrt{1^2 - 4^2 + 6^2} = \sqrt{53}$ for \bullet^4.</p> <p>4. \bullet^6 is not available to candidates who simply state the formula $\cos \theta = \frac{\overline{ED} \cdot \overline{EF}}{\overline{ED} \overline{EF}}$.</p> <p>However, $\cos \theta = \frac{16}{\sqrt{53} \times \sqrt{14}}$ and $\sqrt{53} \times \sqrt{14} \times \cos \theta = 16$ are acceptable for \bullet^6.</p> <p>5. Accept correct answers rounded to 54° or 0.9 radians (or 60 gradians).</p> <p>6. Do not penalise the omission or incorrect use of units.</p> <p>7. \bullet^7 is only available as a result of using a valid strategy.</p> <p>8. \bullet^7 is only available for a single angle.</p> <p>9. For a correct answer with no working award 0/4.</p>			
Commonly Observed Responses:			
<p>Candidate A - poor notation</p> $\begin{pmatrix} 1 \\ -4 \\ 6 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} = \begin{pmatrix} 2 \\ -4 \\ 18 \end{pmatrix} = 16$ <p style="text-align: right;">$\bullet^3 \times$</p>	<p>Candidate B - insufficient communication</p> $\overline{ED} = \sqrt{53}$ $\overline{EF} = \sqrt{14}$ $\frac{16}{\sqrt{53} \times \sqrt{14}}$ <p>54.028...° or 0.942... radians</p> <p style="text-align: right;">$\bullet^4 \checkmark$ $\bullet^5 \checkmark$ $\bullet^6 \wedge$ $\bullet^7 \checkmark_1$</p>		
<p>Candidate C - BEWARE</p> $\overline{OF} = \sqrt{14}$ <p style="text-align: right;">$\bullet^5 \times$</p>			

Question		Generic scheme	Illustrative scheme	Max mark
4.	(a)	<ul style="list-style-type: none"> •¹ identify x-coordinate •² identify y-coordinate 	<ul style="list-style-type: none"> •¹ 3 •² 5 	2

Notes:

Commonly Observed Responses:


(b)	<ul style="list-style-type: none"> •³ identify roots •⁴ interpret point of inflection •⁵ identify orientation and complete cubic curve 	<ul style="list-style-type: none"> •³ “cubic” with roots at -1 and 2 •⁴ “cubic” with turning point at $(2,0)$ •⁵ cubic with maximum turning point at $(2,0)$ 	3
-----	--	--	---

Notes:


1. Note that the position of the minimum turning point of $f'(x)$ is not being assessed.
2. Where a candidate has not drawn a cubic curve or their graph does not extend outwith $-1 \leq x \leq 2$ award 0/3. However see Candidate D.
3. Do not penalise the appearance of an additional root outwith $-1 \leq x \leq 2$ (on a cubic curve) at •³.



Commonly Observed Responses:

Candidate A - $-f'(x)$



Candidate B



Question	Generic scheme	Illustrative scheme	Max mark
4. (b) (continued)			
<p data-bbox="124 282 293 315">Candidate C</p>  <p data-bbox="124 616 207 649">✓ x x</p>	<p data-bbox="809 282 1458 353">Candidate D - left derivative \neq right derivative at (2,0)</p>  <p data-bbox="809 638 893 672">✓ ✓ x</p>		

Question		Generic scheme	Illustrative scheme	Max mark
6.		<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ state linear equation •² introduce logs •³ use laws of logs •⁴ use laws of logs •⁵ state a and b 	<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $\log_5 y = 3\log_5 x - 2$ •² $\log_5 y = 3\log_5 x - 2\log_5 5$ •³ $\log_5 y = \log_5 x^3 - \log_5 5^2$ •⁴ $\log_5 y = \log_5 \frac{x^3}{5^2}$ •⁵ $a = \frac{1}{25}, b = 3$ or $y = \frac{x^3}{25}$ 	5
		<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ state linear equation •² use laws of logs •³ use laws of logs •⁴ use laws of logs •⁵ state a and b 	<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $\log_5 y = 3\log_5 x - 2$ •² $\log_5 y = \log_5 x^3 - 2$ •³ $\log_5 \frac{y}{x^3} = -2$ •⁴ $\frac{y}{x^3} = 5^{-2}$ •⁵ $a = \frac{1}{25}, b = 3$ or $y = \frac{x^3}{25}$ 	5
		<p style="text-align: center;">Method 3</p> <ul style="list-style-type: none"> •¹ introduce logs to $y = ax^b$ •² use laws of logs •³ interpret intercept •⁴ use laws of logs •⁵ interpret gradient 	<p style="text-align: center;">Method 3</p> <p style="text-align: center;">The equations at •¹, •² and •³ must be stated explicitly</p> <ul style="list-style-type: none"> •¹ $\log_5 y = \log_5 ax^b$ •² $\log_5 y = b\log_5 x + \log_5 a$ •³ $\log_5 a = -2$ •⁴ $a = \frac{1}{25}$ •⁵ $b = 3$ 	5

Question	Generic scheme	Illustrative scheme	Max mark
6. (continued)			
Notes			
<p>1. In any method, marks may only be awarded within a valid strategy using $y = ax^b$. For example, see Candidates C and D.</p> <p>2. Markers must identify the method which best matches the candidate's approach; markers must not mix and match between methods.</p> <p>3. Penalise the omission of base 5 at most once in any method.</p> <p>4. Where candidates use an incorrect base then only \bullet^2 and \bullet^3 are available (in any method).</p> <p>5. Do not accept $a = 5^{-2}$.</p> <p>6. In Method 3, do not accept $m = 3$ or gradient = 3 for \bullet^5.</p> <p>7. Do not penalise candidates who score out "log" from equations of the form $\log_5 m = \log_5 n$.</p>			
Commonly Observed Responses			
<p>Candidate A - missing equations at \bullet^1, \bullet^2 and \bullet^3 in Method 3</p> $a = \frac{1}{25} \quad \bullet^4 \checkmark$ $b = 3 \quad \bullet^5 \checkmark$	<p>Candidate B - no working - Method 3</p> $b = \frac{1}{25} \quad \bullet^4 \times$ $a = 3 \quad \bullet^5 \times$		
<p>Candidate C - Method 2</p> $y = 3x - 2$ <hr style="width: 10%; margin-left: 0;"/> $\log_5 y = 3 \log_5 x - 2 \quad \bullet^1 \checkmark$ $\log_5 y = \log_5 x^3 - 2 \quad \bullet^2 \checkmark$ $y = x^3 - 2 \quad \bullet^3 \times \quad \bullet^4 \times \quad \bullet^5 \times$	<p>Candidate D - Method 2</p> $\log_5 y = 3 \log_5 x - 2 \quad \bullet^1 \checkmark$ $\log_5 y = \log_5 x^3 - 2 \quad \bullet^2 \checkmark$ $\frac{y}{x^3} = -2 \quad \bullet^3 \times \quad \bullet^4 \times \quad \bullet^5 \times$		
<p>Candidate E - use of coordinate pairs</p> $\log_5 x = 4 \text{ and } \log_5 y = 10 \quad \bullet^1 \checkmark$ $x = 5^4 \text{ and } y = 5^{10} \quad \bullet^2 \checkmark$ $\log_5 x = 0, \log_5 y = -2 \quad \bullet^3 \checkmark$ $\Rightarrow x = 1, y = 5^{-2} \quad \bullet^3 \checkmark$ $5^{-2} = a \times 1^b \Rightarrow a = \frac{1}{25} \quad \bullet^4 \checkmark$ $5^{10} = 5^{-2} \times 5^{4b} \Rightarrow -2 + 4b = 10 \quad \bullet^5 \checkmark$ $\Rightarrow b = 3$ <p style="text-align: center;">Candidates may use (0, -2) for \bullet^1 and \bullet^2 and (4,10) for \bullet^3.</p>			

Question		Generic scheme	Illustrative scheme	Max mark
7.		<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ integrate using ‘upper’ – ‘lower’ •² identify limits •³ integrate •⁴ substitute limits •⁵ evaluate area 	<p style="text-align: center;">Method 1</p> <ul style="list-style-type: none"> •¹ $\int \left((6 + 4x - 2x^2) - (x^3 - 6x^2 + 11x) \right) dx$ •² $\int_0^2 \left((6 + 4x - 2x^2) - (x^3 - 6x^2 + 11x) \right) dx$ •³ $6x - \frac{7}{2}x^2 + \frac{4}{3}x^3 - \frac{1}{4}x^4$ •⁴ $\left(6(2) - \frac{7}{2}(2)^2 + \frac{4}{3}(2)^3 - \frac{1}{4}(2)^4 \right) - 0$ •⁵ $\frac{14}{3}$ (units²) 	5
		<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ know to integrate between appropriate limits for both equations •² integrate both functions •³ substitute limits into both expressions •⁴ evaluate both integrals •⁵ evidence of subtracting areas 	<p style="text-align: center;">Method 2</p> <ul style="list-style-type: none"> •¹ $\int_0^2 \dots dx$ and $\int_0^2 \dots dx$ •² $6x + \frac{4x^2}{2} - \frac{2x^3}{3}$ and $\frac{x^4}{4} - \frac{6x^3}{3} + \frac{11x^2}{2}$ •³ $\left(6(2) + \frac{4(2)^2}{2} - \frac{2(2)^3}{3} \right) - 0$ and $\left(\frac{(2)^4}{4} - \frac{6(2)^3}{3} + \frac{11(2)^2}{2} \right) - 0$ •⁴ $\frac{44}{3}$ and 10 •⁵ $\frac{14}{3}$ (units²) 	

Question	Generic scheme	Illustrative scheme	Max mark
7. (continued)			
Notes:			
<p>1. Correct answer with no working - award 1/5.</p> <p>2. Do not penalise lack of 'dx' at ●¹ in Method 1.</p> <p>3. In Method 1, limits and 'dx' must appear by the ●² stage for ●² to be awarded and in Method 2 by the ●¹ stage for ●¹ to be awarded.</p> <p>4. In Method 1, treat the absence of brackets at ●¹ stage as bad form only if the correct integrand is obtained. See Candidates C and D.</p> <p>5. Where a candidate differentiates one or more terms, or fails to integrate, no further marks are available.</p> <p>6. In Method 1, accept unsimplified expressions such as $6x + \frac{4x^2}{2} - \frac{2x^3}{3} - \frac{x^4}{4} + \frac{6x^3}{3} - \frac{11x^2}{2}$ at ●³.</p> <p>7. Do not penalise the inclusion of '+c'.</p> <p>8. Do not penalise the continued appearance of the integral sign or dx after integrating.</p> <p>9. ●⁵ is not available where solutions include statements such as '$-\frac{14}{3} = \frac{14}{3}$ square units'. See Candidates A and B.</p> <p>10. In Method 1, where a candidate uses an invalid strategy the only marks available are ●³ for integrating a polynomial with at least four terms (of different degree) and ●⁴ for substituting the limits of 0 and 2 into the resulting expression. However, see Candidate E.</p> <p>11. At ●⁴, do not penalise candidates for who reduce powers of 0. For example writing 0 in place of 0⁴. Similarly, do not penalise candidates writing 0 in place of 6(0). However, candidates who write 0³ in place of 0⁴ or 2(0) in place of 6(0) do not gain ●⁴.</p>			
Commonly Observed Responses:			
<p>Candidate A - switched limits</p> $\int_2^0 \left((6+4x-2x^2) - (x^3-6x^2+11x) \right) dx$ <p style="text-align: right;">●² ✓</p> $= 6x - \frac{7}{2}x^2 + \frac{4}{3}x^3 - \frac{1}{4}x^4$ <p style="text-align: right;">●³ ✓</p> $= 0 - \left(6(2) - \frac{7}{2}(2)^2 + \frac{4}{3}(2)^3 - \frac{1}{4}(2)^4 \right)$ <p style="text-align: right;">●⁴ ✓</p> $= -\frac{14}{3}$ $= \frac{14}{3}$ <p style="text-align: right;">●¹ ✗ ●⁵ ✗</p>		<p>Candidate B - 'lower' - 'upper'</p> $\int_0^2 \left((x^3-6x^2+11x) - (6+4x-2x^2) \right) dx$ <p style="text-align: right;">●² ✓</p> $\int_0^2 x^3 - 4x^2 + 7x - 6 dx$ $= \frac{1}{4}x^4 - \frac{4}{3}x^3 + \frac{7}{2}x^2 - 6x$ <p style="text-align: right;">●³ ✓</p> $\left(\frac{1}{4}(2)^4 - \frac{4}{3}(2)^3 + \frac{7}{2}(2)^2 - 6(2) \right) - (0)$ <p style="text-align: right;">●⁴ ✓</p> $= -\frac{14}{3}$ $\therefore \text{Area} = \frac{14}{3}$ <p style="text-align: right;">●¹ ✓ ●⁵ ✓</p>	

Question	Generic scheme	Illustrative scheme	Max mark
7. (continued)			
<p>Candidate C - missing brackets</p> $\int_0^2 6 + 4x - 2x^2 - x^3 - 6x^2 + 11x \, dx$ $\int_0^2 6 - 7x + 4x^2 - x^3 \, dx$	<p>Candidate D - missing brackets</p> $\int_0^2 6 + 4x - 2x^2 - x^3 - 6x^2 + 11x \, dx$ $\int_0^2 6 + 15x - 8x^2 - x^3 \, dx$ $6x + \frac{15}{2}x^2 - \frac{8}{3}x^3 - \frac{1}{4}x^4$ $\left(6(2) + \frac{15}{2}(2)^2 - \frac{8}{3}(2)^3 - \frac{1}{4}(2)^4\right) - (0)$ $\frac{50}{3}$	<p>•¹ ✗ •² ✓₁</p> <p>•¹ ✓ •² ✓</p> <p>•³ ✓₁</p> <p>•⁴ ✓₁</p> <p>•⁵ ✓₁</p>	
<p>Candidate E - 'upper' + 'lower'</p> $\int_0^2 \left((6 + 4x - 2x^2) + (x^3 - 6x^2 + 11x) \right) dx$ $6x + \frac{15}{2}x^2 - \frac{8}{3}x^3 + \frac{1}{4}x^4$ $\left(6(2) + \frac{15}{2}(2)^2 - \frac{8}{3}(2)^3 + \frac{1}{4}(2)^4\right) - 0$ $\frac{74}{3}$	<p>Candidate F - incorrect substitution</p> $\int_0^2 \left((6 + 4x - 2x^2) - (x^3 - 6x^2 + 11x) \right) dx$ $\left(6x + 2x^2 - \frac{2}{3}x^3\right) - \left(\frac{1}{4}x^4 - 2x^3 + \frac{11}{2}x^2\right)$ $\left(6(2) + 2(2)^2 - \frac{2}{3}(2)^3\right) - \left(\frac{1}{4}(0)^4 - 2(0)^3 + \frac{11}{2}(0)^2\right)$ $\frac{44}{3}$	<p>•¹ ✗ •² ✓₁</p> <p>•¹ ✓ •² ✓</p> <p>•³ ✓</p> <p>•⁴ ✓₁</p> <p>•⁴ ✗</p> <p>•⁵ ✓₁</p> <p>•⁵ ✓₂</p>	

Question		Generic scheme	Illustrative scheme	Max mark
8.	(a)	<ul style="list-style-type: none"> •¹ interpret notation •² state expression for $f(g(x))$ 	<ul style="list-style-type: none"> •¹ $f(x+1)$ or $2g(x)^2 - 18$ •² $2(x+1)^2 - 18$ 	2
Notes:				
1. For $2(x+1)^2 - 18$ without working, award both • ¹ and • ² .				
Commonly Observed Responses:				
Candidate A - $g(f(x))$			Candidate B - beware of two "attempts"	
$2x^2 - 17$		• ¹ ✗ • ² ✓ ₁	$f(g(x)) = 2x^2 - 18$ • ¹ ✗ • ² ✗ $f(x+1) = 2(x+1)^2 - 18$	
	(b)	<ul style="list-style-type: none"> •³ apply condition •⁴ state values of x 	<ul style="list-style-type: none"> •³ $2(x+1)^2 - 18 = 0$ •⁴ -4 and 2 	2
Notes:				
2. Working at • ³ must be consistent with working at • ² .				
3. Accept $2(x+1)^2 - 18 \neq 0$ for • ³ only when $x = -4$ and $x = 2$ are stated explicitly at • ⁴ . See Candidate H				
4. • ⁴ is only available for finding the roots of a quadratic.				
5. For subsequent incorrect working, the final mark is not available. For example $-4 < x < 2$.				
Commonly Observed Responses:				
Candidate C - expanding brackets in (a)			Candidate D - expanding brackets in (a)	
Part (a)			Part (a)	
$f(g(x)) = 2(x+1)^2 - 18$		• ¹ ✓ • ² ✓	$f(g(x)) = 2(x+1)^2 - 18$	
$f(g(x)) = 2x^2 + 4x - 16$			$f(g(x)) = 2x^2 - 16$	
Part (b)			Part (b)	
$2x^2 + 4x - 16 = 0$		• ³ ✓	$2x^2 - 16 = 0$	
$x = -4$ and $x = 2$		• ⁴ ✓	$x = \pm\sqrt{8}$	
Candidate E - $g(f(x))$			Candidate F - equivalent condition	
Part (a)				
$f(g(x)) = 2x^2 - 17$		• ¹ ✗ • ² ✓ ₁	$2(x+1)^2 = 18$	
Part (b)				
$2x^2 - 17 = 0$		• ³ ✓ ₁		
$x = \pm\sqrt{\frac{17}{2}}$		• ⁴ ✓ ₁		
Candidate G - use of \neq			Candidate H - use of \neq	
$2(x+1)^2 - 18 \neq 0$		• ³ ✗	$2(x+1)^2 - 18 \neq 0$	
$x \neq -4, x \neq 2$		• ⁴ ✓ ₁	$x \neq -4, x \neq 2$	
			$x = -4, x = 2$	
			• ³ ✓	
			• ⁴ ✓	

Question		Generic scheme	Illustrative scheme	Max mark
9.	(a)	<ul style="list-style-type: none"> •¹ differentiate two non-constant terms •² complete derivative and equate to 0 •³ find x-coordinates •⁴ find y-coordinates 	<ul style="list-style-type: none"> •¹ eg $x^2 - 2x$ •² $x^2 - 2x - 3 = 0$ •³ $-1, 3$ •⁴ $\frac{8}{3}, -8$ 	4

Notes:

1. For a numerical approach, award 0/4.
2. •² is only available if ‘= 0’ appears at the •² stage or in working leading to •³. However, see Candidate A.
3. •³ is only available for solving a quadratic equation.
4. •³ and •⁴ may be awarded vertically.

Commonly Observed Responses:

Candidate A

Stationary points when $\frac{dy}{dx} = 0$

$\frac{dy}{dx} = x^2 - 2x - 3$ •¹ ✓ •² ✓

$\frac{dy}{dx} = (x+1)(x-3)$

$x = -1, 3$ •³ ✓

$y = \frac{8}{3}, -8$ •⁴ ✓

Candidate B - derivative never equated to 0

$x^2 - 2x - 3$ •¹ ✓ •² ^

$(x+1)(x-3)$

$x = -1, 3$ •³ ✓₁

$y = \frac{8}{3}, -8$ •⁴ ✓

	(b)	<ul style="list-style-type: none"> •⁵ evaluate y at $x = 6$ •⁶ state greatest and least values 	<ul style="list-style-type: none"> •⁵ 19 •⁶ greatest = 19 and least = -8 	2
--	-----	--	--	---

Notes:

5. ‘Greatest (6, 19); least (3, -8)’ does not gain •⁶.
6. Where $x = -1$ was not identified as a stationary point in part (a), y must also be evaluated at $x = -1$ to gain •⁶.
7. •⁶ is not available for using y at a value of x , obtained at •³ stage, which lies outwith the interval $-1 \leq x \leq 6$.
8. •⁶ is only available where candidates have attempted to evaluate y at $x = 6$.

Commonly Observed Responses:

--	--	--	--	--

Question		Generic scheme	Illustrative scheme	Max mark
11.	(a)	• ¹ state number of vehicles	• ¹ 6.8 million	1
Notes:				
1. Accept 6.8 or $N = 6.8$ million for • ¹ .				
Commonly Observed Responses:				
	(b)	• ² substitute for N and t • ³ process equation • ⁴ express in logarithmic form • ⁵ solve for k	• ² $125 = 6.8e^{10k}$ stated or implied by • ³ • ³ $\frac{125}{6.8} = e^{10k}$ • ⁴ $\log_e \left(\frac{125}{6.8} \right) = 10k$ • ⁵ 0.2911...	4
Notes:				
2. Accept answers which round to 0.29. 3. Do not penalise rounding or transcription errors (which are correct to 2 significant figures) in intermediate calculations. 4. • ³ may be assumed by • ⁴ . 5. Any base may be used at • ⁴ stage. See Candidate A. 6. At • ⁴ all exponentials must be processed. 7. Accept $\log_e \frac{125}{6.8} = 10k \log_e e$ for • ⁴ . 8. The calculation at • ⁵ must follow from the valid use of exponentials and logarithms at • ³ and • ⁴ . 9. For candidates with no working, or who adopt an iterative approach to arrive at $k = 0.29$, award 1/4. However, if, in the iterations N is calculated for $k = 0.295$ and $k = 0.285$, then award 4/4.				
Commonly Observed Responses:				
Candidate A - use of alternative base $125 = 6.8e^{10k}$ • ² ✓ $\frac{125}{6.8} = e^{10k}$ • ³ ✓ $\log_{10} \left(\frac{125}{6.8} \right) = 10k \log_{10} e$ • ⁴ ✓ $k = 0.2911...$ • ⁵ ✓		Candidate B - missing lines of working $125 = 6.8e^{10k}$ • ² ✓ $k = 0.2911...$ • ³ ^ • ⁴ ^ • ⁵ ✓		
Candidate C - errors in substitution $125000000 = 6.8e^{10k}$ • ² ✗ $\frac{125000000}{6.8} = e^{10k}$ • ³ ✓ ₁ $16.726... = 10k$ • ⁴ ✓ ₁ $k = 1.6726...$ • ⁵ ✓ ₁				

Question	Generic scheme	Illustrative scheme	Max mark
12.	<ul style="list-style-type: none"> •¹ substitute appropriate double angle formula •² factorise •³ solve for $\tan x^\circ$ •⁴ solve $\tan x^\circ = 4$ •⁵ solve $\sin x^\circ = 0$ 	<ul style="list-style-type: none"> •¹ $2(2 \sin x^\circ \cos x^\circ) - \sin^2 x^\circ (= 0)$ •² $\sin x^\circ(4 \cos x^\circ - \sin x^\circ) = 0$ •³ $\tan x^\circ = 4$ (since $x = 90, 270$ are not solutions) •⁴ 76, 256 •⁵ 0, 180 	5

Notes:

1. •¹ is still available to candidates who correctly substitute for $\sin^2 x$ in addition to $\sin 2x$.
2. Substituting $2 \sin A \cos A$ for $\sin 2x^\circ$ at the •¹ stage should be treated as bad form provided the equation is written in terms of x at the •² stage. Otherwise, •¹ is not available.
3. '= 0' must appear by the •² stage for •² to be awarded.
4. Award •² for ' $S(4C - S) = 0$ ' only where $\sin x^\circ = 0$ and $4 \cos x^\circ - \sin x^\circ = 0$ appear.
5. Do not penalise the omission of degree signs.
6. At •³ stage, candidates are not required to check that 90 and 270 are not solutions before dividing by $\cos x^\circ$. Where candidates have divided by $\sin x$ at the •² stage without considering $\sin x = 0$, •³ and •⁴ are still available.
7. At •³ stage, candidates may use the wave function and arrive at $\sqrt{17} \cos(x + 14)^\circ = 0$, or an equivalent wave form, instead of $\tan x^\circ = 4$.
8. •⁴ is only available where the working at the •³ stage is of equivalent difficulty to reaching $\tan x^\circ = 4$.
9. •⁵ is not available where $\sin x = 0$ comes from an invalid strategy.
10. For candidates who work only in radians, •⁵ is not available.
11. •⁴ and •⁵ may be awarded vertically. See also Candidate B.
12. Do not penalise solutions outwith $0 \leq x < 360$.

Commonly Observed Responses:

<p>Candidate A - working in radians</p> <p>∴</p> <p>$\tan x^\circ = 4$</p> <p>1.326, 4.468</p> <p>0, π</p>	<p>•¹ ✓ •² ✓</p> <p>•³ ✓</p> <p>•⁴ ✓₁</p> <p>•⁵ ✓₂</p>	<p>Candidate B - partial solutions</p> <p>$2(2 \sin x^\circ \cos x^\circ) - \sin^2 x^\circ = 0$</p> <p>$\sin x^\circ(4 \cos x^\circ - \sin x^\circ) = 0$</p> <p>$\sin x^\circ = 0$</p> <p>$x = 180$</p> <p>$\tan x^\circ = 4$</p> <p>$x = 76$</p> <p>•⁵ ^</p>	<p>•¹ ✓</p> <p>•² ✓</p> <p>•³ ✓</p> <p>•⁴ ✓</p>
---	---	---	---

Question		Generic scheme	Illustrative scheme	Max mark
13.		<ul style="list-style-type: none"> •¹ state repeated factor •² state non-repeated linear factors •³ calculate k and express in required form 	<ul style="list-style-type: none"> •¹ $(x-3)^2(\dots)(\dots)$ •² $(\dots)^2(x+1)(x-5)$ •³ $f(x) = \frac{1}{5}(x-3)^2(x+1)(x-5)$ 	3
Notes:				
1. Do not penalise the omission of $f(x) =$ or the inclusion of $y =$.				
2. Accept $f(x) = \frac{1}{5}(x+3)^2(x+1)(x+5)$ for • ³ .				
Commonly Observed Responses:				
Candidate A - incorrect signs		Candidate B - incorrect repeated root		
$f(x) = k(x+3)^2(x-1)(x+5)$		$f(x) = k(x+1)^2(x-3)(x-5)$		
• ¹ ✗ • ² ✓ ₁		• ¹ ✗ • ² ✓ ₁		
$f(x) = \frac{1}{5}(x+3)^2(x-1)(x+5)$		$f(x) = -\frac{3}{5}(x+1)^2(x-3)(x-5)$		
• ³ ✓ ₁		• ³ ✓ ₁		
Candidate C - incorrect repeated root		Candidate D - incorrect signs and repeated root		
$f(x) = k(x-5)^2(x+1)(x-3)$		$f(x) = k(x+5)^2(x-1)(x+3)$		
• ¹ ✗ • ² ✓ ₁		• ¹ ✗ • ² ✗		
$f(x) = \frac{3}{25}(x-5)^2(x+1)(x-3)$		$f(x) = \frac{3}{25}(x+5)^2(x-1)(x+3)$		
• ³ ✓ ₁		• ³ ✓ ₁		
Candidate E - incorrect signs and repeated root		Candidate F - use of a, b and c		
$f(x) = k(x-1)^2(x+5)(x+3)$		$a = -3$		
• ¹ ✗ • ² ✗		$b = 1, c = -5$ (or $b = -5, c = 1$)		
$f(x) = -\frac{3}{5}(x-1)^2(x+5)(x+3)$		$k = \frac{1}{5}$		
• ³ ✓ ₁		• ¹ ✓		
		• ² ✓		
		• ³ ^		

[END OF MARKING INSTRUCTIONS]

General marking principles for Higher Mathematics

Always apply these general principles. Use them in conjunction with the detailed marking instructions, which identify the key features required in candidates' responses.

For each question, the marking instructions are generally in two sections:

- *generic scheme* – this indicates why each mark is awarded
- *illustrative scheme* – this covers methods which are commonly seen throughout the marking

In general, you should use the illustrative scheme. Only use the generic scheme where a candidate has used a method not covered in the illustrative scheme.

- Always use positive marking. This means candidates accumulate marks for the demonstration of relevant skills, knowledge and understanding; marks are not deducted for errors or omissions.
- If you are uncertain how to assess a specific candidate response because it is not covered by the general marking principles or the detailed marking instructions, you must seek guidance from your team leader.
- One mark is available for each •. There are no half marks.
- If a candidate's response contains an error, all working subsequent to this error must still be marked. Only award marks if the level of difficulty in their working is similar to the level of difficulty in the illustrative scheme.
- Only award full marks where the solution contains appropriate working. A correct answer with no working receives no mark, unless specifically mentioned in the marking instructions.
- Candidates may use any mathematically correct method to answer questions, except in cases where a particular method is specified or excluded.
- If an error is trivial, casual or insignificant, for example $6 \times 6 = 12$, candidates lose the opportunity to gain a mark, except for instances such as the second example in point (h) below.
- If a candidate makes a transcription error (question paper to script or within script), they lose the opportunity to gain the next process mark, for example

This is a transcription error and so the mark is not awarded.

This is no longer a solution of a quadratic equation, so the mark is not awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$x = 1$$

The following example is an exception to the above

This error is not treated as a transcription error, as the candidate deals with the intended quadratic equation. The candidate has been given the benefit of the doubt and all marks awarded.

$$x^2 + 5x + 7 = 9x + 4$$

$$x - 4x + 3 = 0$$

$$(x - 3)(x - 1) = 0$$

$$x = 1 \text{ or } 3$$

(i) **Horizontal/vertical marking**

If a question results in two pairs of solutions, apply the following technique, but only if indicated in the detailed marking instructions for the question.

Example:

$$\begin{array}{cc} \bullet^5 & \bullet^6 \\ \bullet^5 & x = 2 \quad x = -4 \\ \bullet^6 & y = 5 \quad y = -7 \end{array}$$

Horizontal: $\bullet^5 x = 2$ and $x = -4$ Vertical: $\bullet^5 x = 2$ and $y = 5$
 $\bullet^6 y = 5$ and $y = -7$ $\bullet^6 x = -4$ and $y = -7$

You must choose whichever method benefits the candidate, **not** a combination of both.

(j) In final answers, candidates should simplify numerical values as far as possible unless specifically mentioned in the detailed marking instruction. For example

$$\begin{array}{ll} \frac{15}{12} \text{ must be simplified to } \frac{5}{4} \text{ or } 1\frac{1}{4} & \frac{43}{1} \text{ must be simplified to } 43 \\ \frac{15}{0.3} \text{ must be simplified to } 50 & \frac{4}{3} \text{ must be simplified to } \frac{4}{15} \\ \sqrt{64} \text{ must be simplified to } 8^* & \end{array}$$

*The square root of perfect squares up to and including 144 must be known.

(k) Commonly Observed Responses (CORs) are shown in the marking instructions to help mark common and/or non-routine solutions. CORs may also be used as a guide when marking similar non-routine candidate responses.

(l) Do not penalise candidates for any of the following, unless specifically mentioned in the detailed marking instructions:

- working subsequent to a correct answer
- correct working in the wrong part of a question
- legitimate variations in numerical answers/algebraic expressions, for example angles in degrees rounded to nearest degree
- omission of units
- bad form (bad form only becomes bad form if subsequent working is correct), for example $(x^3 + 2x^2 + 3x + 2)(2x + 1)$ written as $(x^3 + 2x^2 + 3x + 2) \times 2x + 1$
 $= 2x^4 + 5x^3 + 8x^2 + 7x + 2$ gains full credit
- repeated error within a question, but not between questions or papers

(m) In any 'Show that...' question, where candidates have to arrive at a required result, the last mark is not awarded as a follow-through from a previous error, unless specified in the detailed marking instructions.

- (n) You must check all working carefully, even where a fundamental misunderstanding is apparent early in a candidate's response. You may still be able to award marks later in the question so you must refer continually to the marking instructions. The appearance of the correct answer does not necessarily indicate that you can award all the available marks to a candidate.
- (o) You should mark legible scored-out working that has not been replaced. However, if the scored-out working has been replaced, you must only mark the replacement working.
- (p) If candidates make multiple attempts using the same strategy and do not identify their final answer, mark all attempts and award the lowest mark. If candidates try different valid strategies, apply the above rule to attempts within each strategy and then award the highest mark.

For example:

Strategy 1 attempt 1 is worth 3 marks.	Strategy 2 attempt 1 is worth 1 mark.
Strategy 1 attempt 2 is worth 4 marks.	Strategy 2 attempt 2 is worth 5 marks.
From the attempts using strategy 1, the resultant mark would be 3.	From the attempts using strategy 2, the resultant mark would be 1.

In this case, award 3 marks.